

Involutions of Iwahori-Hecke algebras and representations of fixed subalgebras

Hideo Mitsuhashi

Department of Mechanical Engineering

Tomakomai National College of Technology

443 Nishikioka, Tomakomai, Hokkaido, 059-1275, Japan

Abstract

We establish branching rules between some Iwahori-Hecke algebra of type B and their subalgebras which are defined as fixed subalgebras by involutions including Goldman involution. The Iwahori-Hecke algebra of type D is one of such fixed subalgebras. We also obtain branching rules between those fixed subalgebras and their intersection subalgebra. We determine basic sets of irreducible representations of those fixed subalgebras and their intersection subalgebra by making use of generalized Clifford theory.

1 Introduction

In our paper [5], we gave the q -analogue of the alternating group and its irreducible representations. Since the Iwahori-Hecke algebra of type A can be considered as a q -analogue of the symmetric group, we defined the q -analogue of the alternating group as a subalgebra of it. An extension of [5] to the Hecke algebra of type B was given in [6]. We showed that those subalgebras can be defined as fixed subalgebras by Goldman involutions of Hecke algebras in [6]. The Iwahori-Hecke algebra of type D can be considered as a subalgebra of some specialized Iwahori-Hecke algebra of type B , and can be realized as a fixed subalgebra by an involution, which we denote by $\sharp$, of the specialized Iwahori-Hecke algebra. The specialized Iwahori-Hecke algebra has a $\mathbb{Z}_2$ -graded Clifford system whose submodule including identity element coincides with the Iwahori-Hecke algebra of type D . Using generalized Clifford theory, we give a new proof of the branching rule between the specialized Iwahori-Hecke algebra and the Iwahori-Hecke algebra of type D , which has already been obtained in [4]. We do not specify the base fields in the statements below in this section because of simplicity. The details are specified in corresponding paragraphs since the next chapter. Let π_λ be the irreducible representation of the Iwahori-Hecke algebra of type B corresponding to $\lambda = (\lambda^{(1)}, \lambda^{(2)})$, a 2-tuple of Young diagrams. Then we have the following.

Theorem 6.8. ($\sharp 1$) The restrictions $\pi_{\lambda, \sharp}$ and $\pi_{\lambda^*, \sharp}$ of irreducible representations π_λ and π_{λ^*} corresponding to $\lambda = (\lambda^{(1)}, \lambda^{(2)})$ and $\lambda^* = (\lambda^{(2)}, \lambda^{(1)})$ to the fixed subalgebra by $\sharp$ are equivalent.

($\sharp 2$) If $\lambda^* = \lambda$, then $\pi_{\lambda, \sharp}$ decomposes into two inequivalent subrepresentations $\pi_{\lambda, \sharp}^+$ and $\pi_{\lambda, \sharp}^-$ of the same degree.

($\sharp 3$) Let $\{\lambda_i, \lambda_i^*, \mu_j\}_{i,j}$ be the set of 2-tuples of Young diagrams such that $\lambda_i \neq \lambda_i^*$ and $\mu_j = \mu_j^*$. Then $\{\pi_{\lambda_i, \sharp}, \pi_{\mu_j, \sharp}^+, \pi_{\mu_j, \sharp}^-\}_{i,j}$ is a basic set of irreducible representations of the fixed subalgebra by $\sharp$.

Besides such involution, there are two other involutions of such specialized Iwahori-Hecke algebra. One of them is Goldman involution, which we denote by $\natural$. Goldman involution itself can be defined for generic (non-specialized) Iwahori-Hecke algebras of type B . We have already given the fixed subalgebra by Goldman involution in generic Iwahori-Hecke algebras of type B , and have determined the branching rule and a basic set of irreducible representations in [6] as follows.

Theorem 6.2. ($\natural 1$) The restrictions $\pi_{\lambda, \natural}$ and $\pi_{\lambda'^*, \natural}$ of irreducible representations π_λ and $\pi_{\lambda'^*}$ corresponding to $\lambda = (\lambda^{(1)}, \lambda^{(2)})$ and $\lambda'^* = (\lambda^{(2)'}, \lambda^{(1)'})$ to the fixed subalgebra by $\natural$ are equivalent, where $\lambda^{(k)'} stands for the transpose of $\lambda^{(k)}$.$

($\natural 2$) If $\lambda'^* = \lambda$, then $\pi_{\lambda, \natural}$ decomposes into two inequivalent subrepresentations $\pi_{\lambda, \natural}^+$ and $\pi_{\lambda, \natural}^-$ of the same degree.

($\natural 3$) Let $\{\lambda_i, \lambda_i', \mu_j\}_{i,j}$ be the set of 2-tuples of Young diagrams such that $\lambda_i \neq \lambda_i'$ and $\mu_j = \mu_j'$. Then $\{\pi_{\lambda_i, \natural}, \pi_{\mu_j, \natural}^+, \pi_{\mu_j, \natural}^-\}_{i,j}$ is a basic set of irreducible representations of the fixed subalgebra by $\natural$.

The other, which we denote by $\flat$, is new. We give the fixed subalgebra by it and determine properties of it. In the same manner as in the case $\sharp$ or $\natural$, we obtain the branching rule and a basic set of irreducible representations of the fixed subalgebra.

Theorem 6.13. ($\flat 1$) The restrictions $\pi_{\lambda, \flat}$ and $\pi_{\lambda', \flat}$ of irreducible representations π_λ and $\pi_{\lambda'}$ corresponding to $\lambda = (\lambda^{(1)}, \lambda^{(2)})$ and $\lambda' = (\lambda^{(1)'}, \lambda^{(2)'})$ to the fixed subalgebra by $\flat$ are equivalent.

($\flat 2$) If $\lambda' = \lambda$, then $\pi_{\lambda, \flat}$ decomposes into two inequivalent subrepresentations $\pi_{\lambda, \flat}^+$ and $\pi_{\lambda, \flat}^-$ of the same degree.

($\flat 3$) Let $\{\lambda_i, \lambda_i', \mu_j\}_{i,j}$ be the set of 2-tuples of Young diagrams such that $\lambda_i \neq \lambda_i'$ and $\mu_j = \mu_j'$. Then $\{\pi_{\lambda_i, \flat}, \pi_{\mu_j, \flat}^+, \pi_{\mu_j, \flat}^-\}_{i,j}$ is a basic set of irreducible representations of the fixed subalgebra by $\flat$.

The intersection of those three fixed subalgebras can be considered as the fixed subalgebra by Goldman involution in the Iwahori-Hecke algebra of type D , while it may be viewed as a subalgebra of both the fixed subalgebra by $\natural$ and that by $\flat$. We denote by $\pi_{\lambda, \dagger}, \pi_{\lambda, \natural, \dagger}, \pi_{\lambda, \sharp, \dagger}, \pi_{\lambda, \flat, \dagger}$ the restrictions of $\pi_\lambda, \pi_{\lambda, \natural}, \pi_{\lambda, \sharp}, \pi_{\lambda, \flat}$ to the intersection respectively. It is clear that $\pi_{\lambda, \dagger} \cong \pi_{\lambda, \natural, \dagger} \cong \pi_{\lambda, \sharp, \dagger} \cong \pi_{\lambda, \flat, \dagger}$. We give branching rules between those fixed subalgebras and the intersection subalgebra of them, and determine a basic set of irreducible representations of the intersection subalgebra as follows.

(†1) (Proposition 7.1) $\pi_{\lambda, \dagger} \cong \pi_{\lambda^*, \dagger} \cong \pi_{\lambda', \dagger} \cong \pi_{\lambda'^*, \dagger}$.

(†2) (Proposition 7.3)

- (a) If $\lambda^* = \lambda$, $\lambda' \neq \lambda$, $\lambda'^* \neq \lambda$, then $\pi_{\lambda, \#}^+ \cong \pi_{\lambda', \#}^+$ and $\pi_{\lambda, \#}^- \cong \pi_{\lambda', \#}^-$. They are irreducible and mutually inequivalent.
- (b) If $\lambda^* \neq \lambda$, $\lambda' = \lambda$, $\lambda'^* \neq \lambda$, then $\pi_{\lambda, b}^+ \cong \pi_{\lambda', b}^+$ and $\pi_{\lambda, b}^- \cong \pi_{\lambda', b}^-$. They are irreducible and mutually inequivalent.
- (c) If $\lambda^* \neq \lambda$, $\lambda' \neq \lambda$, $\lambda'^* = \lambda$, then $\pi_{\lambda, \natural}^+ \cong \pi_{\lambda', \natural}^+$ and $\pi_{\lambda, \natural}^- \cong \pi_{\lambda', \natural}^-$. They are irreducible and mutually inequivalent.

(†3) (Corollary 7.7) If $\lambda^* = \lambda' = \lambda'^* = \lambda$, then $\pi_{\lambda, \dagger}$ decomposes into four inequivalent subrepresentations $\pi_{\lambda, \dagger}^{++}$, $\pi_{\lambda, \dagger}^{+-}$, $\pi_{\lambda, \dagger}^{-+}$ and $\pi_{\lambda, \dagger}^{--}$ of the same degree.

(†4) (Theorem 7.9) Let $\{\lambda_i, \lambda_i^*, \lambda'_i, \lambda'^*_i, \mu_j, \mu_j', \nu_k, \nu_k^*, \kappa_l, \kappa_l', \iota_m\}_{i,j,k,l,m}$ be the set of 2-tuples of Young diagrams such that

$$\begin{aligned} \lambda_i' &\neq \lambda_i, \lambda_i^* \neq \lambda_i, \lambda'^*_i \neq \lambda_i, & \mu_j^* &= \mu_j, \mu_j' \neq \mu_j, \mu'^*_j \neq \mu_j, \\ \nu_k^* &\neq \nu_k, \nu_k' = \nu_k, \nu_k'^* \neq \nu_k, & \kappa_l^* &\neq \kappa_l, \kappa_l' \neq \kappa_l, \kappa_l'^* = \kappa_l, & \iota_m' &= \iota_m^* = \iota'^*_m = \iota_m. \end{aligned}$$

Then

$$\{\pi_{\lambda_i, \dagger}, \pi_{\mu_j, \#}^\pm, \pi_{\nu_k, b}^\pm, \pi_{\kappa_l, \natural}^\pm, \pi_{\iota_m, \dagger}^{++}, \pi_{\iota_m, \dagger}^{+-}, \pi_{\iota_m, \dagger}^{-+}, \pi_{\iota_m, \dagger}^{--}\}_{i,j,k,l,m}$$

is a basic set of irreducible representations of the intersection subalgebra.

2 Preliminaries on group graded algebras

In this section, we review some properties about group graded algebras. Statements listed below and its proof can be found in Curtis-Reiner [2]. For a detailed account of group graded rings, we refer to [7]. Here we give the definition of the S -graded Clifford system in the algebra over the commutative ring for the finite group S and give properties of S -graded Clifford systems. The definition of the S -graded R -algebra given in [2] is somewhat unpopular, so we adopt the term “ S -crossed product” instead of “ S -graded R -algebra” in accordance with [7].

Definition 2.1. Let S be a finite group, R a commutative ring, and A an R -algebra, finitely generated over R as module. A family of R -submodules $\{A_s\}_{s \in S}$ of A , indexed by the elements of S , is called an S -graded Clifford system in A , if the following conditions are satisfied.

(C1) $A_s A_t = A_{st}$

(C2) For each $s \in S$, there exists a unit $a_s \in A$ such that $A_s = a_s A_1 = A_1 a_s$

(C3) $A = \bigoplus_{s \in S} A_s$.

$$(C4) \quad 1 \in A_1$$

An R -algebra satisfying these conditions is called an S -crossed product. We notice that the S -graded Clifford system is a generalization of the twisted group algebra $R[S]_\alpha$ with factor set α , and trivial G -action on R . The next five results are basic facts about the representation theory of the S -graded Clifford system. We will use these results to derive the branching rule later. The proofs of these results are shown in Curtis-Reiner [2] Chapter 1, Section 11C (p.267-279).

Proposition 2.2 ([2] (11.13)). *Let A be an R -algebra with an S -graded Clifford system, and let L be a left A_1 -module, and M a left A -module. Then there are two isomorphisms of R -modules*

$$(1) \quad \text{Hom}_{A_1}(L, M_{A_1}) \cong \text{Hom}_A(L^A, M)$$

and

$$(2) \quad \text{Hom}_{A_1}(M_{A_1}, L) \cong \text{Hom}_A(M, L^A),$$

where M_{A_1} is the A_1 -module obtained by restriction of scalars from A to A_1 , and L^A is the induced module from A_1 of A defined as follows.

$$L^A = A \otimes_{A_1} L$$

A left A_1 -module L is called *stable relative to A* if L is A_1 -isomorphic to all $a_s \otimes L$ with $s \in S$.

Proposition 2.3 ([2] (11.14)). *Let A have an S -graded Clifford system, let L be a left A_1 -module, and let E denote the endomorphism algebra $\text{End}_A(L^A)$, viewed as a ring of right operators on L^A . For each $s \in S$, let*

$$E_s = \{f \in E \mid (1 \otimes L)f \subseteq a_s \otimes L\}$$

Then

(1) *For all $s, t \in S$, we have*

$$A_s(a_t \otimes L) = a_{st} \otimes L, \quad (a_s \otimes L)E_s \subseteq (a_{st} \otimes L)$$

$$E_s E_t \subseteq E_{st}, \quad 1 \in E_1, \quad E = \bigoplus_{s \in S} E_s$$

(2) *Each element $\varphi \in \text{Hom}_{A_1}(1 \otimes L, a_s \otimes L)$ extends to a unique element $\hat{\varphi} \in E_s$, given by $(a \otimes l)\hat{\varphi} = a(1 \otimes l)\varphi$ for $l \in L, a \in A$. The map $\varphi \rightarrow \hat{\varphi}$ defines an isomorphism of R -modules*

$$\text{Hom}_{A_1}(1 \otimes L, a_s \otimes L) \cong E_s \quad s \in S$$

and this is an isomorphism of R -algebras when $a_s = 1$.

(3) *If L is stable and R is noetherian, then E has an S -graded Clifford system, with units $\hat{e}_s \in E_s$ defined by (2) from A_1 -isomorphisms $e_s : 1 \otimes L \rightarrow a_s \otimes L$, for all $s \in S$.*

Proposition 2.4 ([2] (11.15)). *Let A have an S -graded Clifford system over a field K , and let L be a simple A_1 -module. Then E_1 is a finite dimensional division algebra over K , and*

$$T = \{t \in S \mid 1 \otimes L \cong a_t \otimes L \text{ as } A_1\text{-modules}\}$$

is a subgroup of S . Further the K -algebra

$$E_T = \sum_{t \in T} E_t$$

is a T -crossed product.

The subgroup T defined in Proposition 2.4 is called the *inertial group* of L .

Proposition 2.5 ([2] (11.16)). *Let A be an S -crossed product over a field K , M a simple A -module, and L a simple submodule of M_{A_1} . Then the following statements hold.*

- (1) M_{A_1} is a semisimple A_1 -module, whose simple summands are isomorphic to the simple A_1 -modules $\{a_s \otimes L \mid s \in S\}$.
- (2) Let $T = \{t \in S \mid L \cong a_t \otimes L \text{ as } A_1\text{-module}\}$, and let $B = \sum_{t \in T} A_t$. Then the homogeneous component N of M_{A_1} containing L is a B -module, and M is A -isomorphic to the induced module $A \otimes_B L$.
- (3) N is a simple submodule of L^B , and L is stable relative to B .

Theorem 2.6 ([2] (11.17)). *Let A be an S -crossed product over a field K , and let L be a simple A_1 -module which is stable relative to A . Let $E = \text{End}_A(L^A)$ as ring of right operators of L^A .*

- (1) *There is an isomorphism θ from the lattice of left ideals I in E onto the lattice of A -submodules U of L^A , given by*

$$U = L^A \cdot I, \quad I = \{\gamma \in E \mid L^A \gamma \subseteq U\}$$

- (2) *The A -module $L^A \cdot I$ corresponding to I is A -isomorphic to $L \otimes_{E_1} I$, where the action of A on this tensor product is given by*

$$a(l \otimes \gamma) = (al)\hat{e}_s^{-1} \otimes \hat{e}_s \gamma \quad a \in A_s, l \in L, \gamma \in I$$

with E_1 and the units $\{\hat{e}_s\}$ in E_s defined as in Proposition 2.3. Furthermore,

$$\dim_K(L \otimes_{E_1} I) = (\dim_{E_1} L)(\dim_K I)$$

- (3) *The lattice isomorphism θ is functorial, in the sense that E -homomorphisms $f : I \rightarrow I'$ between left ideals of E , correspond bijectively to A -homomorphisms $1 \otimes f : L \otimes_{E_1} I \rightarrow L \otimes_{E_1} I'$.*

If $S = \mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ and K is an algebraic closed field, then we obtain the following theorem which we will use later frequently.

Theorem 2.7. *Let A be a $\mathbb{Z}_2$ -crossed product over an algebraic closed field K , and M a simple left A -module. Then one of the following holds :*

- (1) $M_{A_{\bar{0}}}$ is a simple left $A_{\bar{0}}$ -module. The induced module $(M_{A_{\bar{0}}})^A$ is a direct sum of two mutually non-isomorphic simple left A -modules M^1 and M^2 . One of those is A -isomorphic to M and their restrictions $M_{A_{\bar{0}}}^1$ and $M_{A_{\bar{0}}}^2$ are A_1 -isomorphic to $M_{A_{\bar{0}}}$.
- (2) $M_{A_{\bar{0}}} = L_1 \oplus L_2$ where L_1, L_2 are mutually non-isomorphic simple left $A_{\bar{0}}$ -modules which have same dimension. The induced modules L_1^A and L_2^A are A -isomorphic to M . Both L_1 and L_2 never appear as irreducible constituents of $M'_{A_{\bar{0}}}$, where M' is any simple left A -module which is non-isomorphic to M .

Furthermore, all $M_{A_{\bar{0}}}$ in the case (1) and all L_1 and L_2 in the case (2) are mutually non-isomorphic except for the pairs $(M_{A_{\bar{0}}}^1, M_{A_{\bar{0}}}^2)$ where M is in the case (1).

Proof. Let L be a simple submodule of $M_{A_{\bar{0}}}$ and T the inertial group of L . Since T is a subgroup of $\mathbb{Z}_2$, T must be $\{0\}$ or $\mathbb{Z}_2$.

If $T = \{0\}$, then $a_{\bar{1}} \otimes L \not\cong 1 \otimes L$ as $A_{\bar{0}}$ -modules. From Proposition 2.5 (1) we have

$$M_{A_{\bar{0}}} \cong m_{\bar{0}}(1 \otimes L) \oplus m_{\bar{1}}(a_{\bar{1}} \otimes L)$$

as $A_{\bar{0}}$ -modules where $m_{\bar{0}}$ and $m_{\bar{1}}$ are multiplicities of $1 \otimes L$ and $a_{\bar{1}} \otimes L$ respectively. We also have

$$M_{A_{\bar{0}}} = a_{\bar{1}} M_{A_{\bar{0}}} \cong m_{\bar{0}}(a_{\bar{1}} \otimes L) \oplus m_{\bar{1}}(1 \otimes L)$$

Thus $m_{\bar{0}} = m_{\bar{1}}$, so we may write $M_{A_{\bar{0}}} \cong m\{(1 \otimes L) \oplus (a_{\bar{1}} \otimes L)\}$ for some positive integer m . Hence $\dim_K M = 2m \dim_K L$. On the other hand, $\dim_K M \leq 2 \dim_K L$ since M is an irreducible constituent of $L^A = (1 \otimes L) \oplus (a_{\bar{1}} \otimes L)$. Therefore $m = 1$ and $L^A = M$ holds. Let M' be any simple left A -module which is non-isomorphic to M . If $M'_{A_{\bar{0}}}$ contains L_1 or L_2 as an irreducible constituent, say L_1 is contained, then $(L_1)^A$ is A -isomorphic to both M and M' , but this is contradiction. This corresponds to the case (2).

If $T = \mathbb{Z}_2$, then $a_{\bar{1}} \otimes L \cong 1 \otimes L$ as $A_{\bar{0}}$ -modules and $M_{A_{\bar{0}}} \cong mL$ for some positive integer m . $E = \text{End}_A L^A$ has a $\mathbb{Z}_2$ -graded Clifford system by Proposition 2.3 (3). Hence we may write

$$E = E_{\bar{0}} \otimes E_{\bar{1}} = E_{\bar{0}} \otimes E_{\bar{0}} \hat{e}_{\bar{1}}$$

Moreover, since K is algebraic closed we have $E_{\bar{0}} \cong K$. Multiplying $\hat{e}_{\bar{1}}$ by a suitable scalar, we may assume $\hat{e}_{\bar{1}}^2 = 1$. Thus E is isomorphic to the quotient algebra $K[X]/(X^2 - 1)$ of the polynomial ring by the correspondence $\hat{e}_{\bar{1}} \mapsto X$. Let I_1, I_2 be ideals of E which are defined by

$$I_1 = \langle 1 + \hat{e}_{\bar{1}} \rangle, \quad I_2 = \langle 1 - \hat{e}_{\bar{1}} \rangle.$$

Then we readily see that $\dim_K I_1 = \dim_K I_2 = 1$ and $E = I_1 \oplus I_2$. Since I_1 and I_2 are mutually non-isomorphic simple E -modules, $L^A \cdot I_1$ and $L^A \cdot I_2$ are mutually non-isomorphic simple A -modules by Theorem 2.6 (1),(3). From Theorem 2.6 (2), we have

$$\begin{aligned}\dim_K L \cdot I_1 &= \dim_K L \times \dim_K I_1 = \dim_K L, \\ \dim_K L \cdot I_2 &= \dim_K L \times \dim_K I_2 = \dim_K L.\end{aligned}$$

Since $\dim_K L^A = 2 \dim_K L$, L^A decomposes into two simple submodules as follows.

$$L^A = L \cdot I_1 \oplus L \cdot I_2$$

M is an irreducible constituent of L^A , hence $M \cong L \cdot I_1$ or $M \cong L \cdot I_2$. In each case, the restriction M_{A_0} is isomorphic to L by Theorem 2.6 (2). This corresponds to the case (1).

For the last statement, we have already shown that $M_{A_0}^1 \cong M_{A_0}^2$ and $L_1 \not\cong L_2$. Other pairs of simple A_1 -modules are mutually non-isomorphic since their induced modules are mutually non-isomorphic. Thus we have completed the proof. $\square$

3 Hecke algebras of type B_n and D_n

Let $(W, S = \{s_1, \dots, s_n\})$ be a Coxeter system of rank n . Let R be a commutative domain with 1, and let $q_i (i = 1, \dots, n)$ be any invertible elements of R such that $q_i = q_j$ if s_i is conjugate to s_j in W . The Iwahori-Hecke algebra $\mathcal{H}_R(W, S)$ is an R -algebra generated by $\{T_{s_i} | s_i \in S\}$ with the relations:

$$\begin{aligned}(\text{H1}) \quad T_{s_i}^2 &= (q_i - 1)T_{s_i} + q_i && \text{if } i = 1, 2, \dots, n, \\ (\text{H2}) \quad (T_{s_i} T_{s_j})^{k_{ij}} &= (T_{s_j} T_{s_i})^{k_{ij}} && \text{if } m_{ij} = 2k_{ij}, \\ (\text{H3}) \quad (T_{s_i} T_{s_j})^{k_{ij}} T_{s_i} &= (T_{s_j} T_{s_i})^{k_{ij}} T_{s_j} && \text{if } m_{ij} = 2k_{ij} + 1,\end{aligned}$$

where m_{ij} is the order of $s_i s_j$ in W . We define $T_w = T_{s_{i_1}} T_{s_{i_2}} \cdots T_{s_{i_k}}$ where $w = s_{i_1} s_{i_2} \cdots s_{i_k}$ is a reduced expression of w . It is known that $\{T_w | w \in W\}$ form a basis of $\mathcal{H}_R(W, S)$ as free R -modules. The relations (H1)–(H3) is equivalent to the following two relations:

$$\begin{aligned}(\text{h1}) \quad T_{s_i} T_w &= T_{s_i w} && \text{if } l(w) < l(s_i w), \\ (\text{h2}) \quad T_{s_i} T_w &= (q_i - 1)T_w + q_i T_{s_i w} && \text{if } l(w) > l(s_i w),\end{aligned}$$

or equivalently,

$$\begin{aligned}(\text{h'1}) \quad T_w T_{s_i} &= T_{ws_i} && \text{if } l(w) < l(ws_i), \\ (\text{h'2}) \quad T_w T_{s_i} &= (q_i - 1)T_w + q_i T_{ws_i} && \text{if } l(w) > l(ws_i),\end{aligned}$$

where $l(w)$ means the length of w .

We set $u = q_1$, $q = q_2 = q_3 = \cdots q_n$. The Hecke algebra $\mathcal{H}_{R,B_n}(u, q)$ of type B_n ($n \geq 2$) is the algebra over R defined by the generators $a_1, a_2, \dots, a_n$ and the following defining relations.

$$(B1) \quad a_1^2 = (u - 1)a_1 + u$$

$$(B2) \quad a_i^2 = (q - 1)a_i + q \quad \text{if } i = 2, 3, \dots, n$$

$$(B3) \quad a_1 a_2 a_1 a_2 = a_2 a_1 a_2 a_1$$

$$(B4) \quad a_i a_{i+1} a_i = a_{i+1} a_i a_{i+1} \quad \text{if } i = 2, 3, \dots, n - 1$$

$$(B5) \quad a_i a_j = a_j a_i \quad \text{if } |i - j| > 1$$

Let $u = 1$ and consider $\mathcal{H}_{R,B_n}(1, q)$. Then (B1) is reduced to (B1') $a_1^2 = 1$. One can readily check that the element $\bar{a}_1 = a_1 a_2 a_1$ satisfies $\bar{a}_1^2 = (q - 1)\bar{a}_1 + q$. For $n \geq 2$, we define $\bar{\mathcal{H}}_{R,B_n}(1, q)$ to be the subalgebra of $\mathcal{H}_{R,B_n}(1, q)$ generated by $\bar{a}_1, \bar{a}_2, \dots, \bar{a}_n$ where $\bar{a}_i = a_i$ for $i > 1$. We also define $\bar{\mathcal{H}}_{R,B_1}(1, q)$ to be the subalgebra of $\mathcal{H}_{R,B_1}(1, q)$ generated by the identity element. We readily see that $\bar{\mathcal{H}}_{R,B_n}(1, q)$ satisfies the following relations.

$$(D1) \quad \bar{a}_i^2 = (q - 1)\bar{a}_i + q \quad \text{if } i = 1, 2, \dots, n$$

$$(D2) \quad \bar{a}_1 \bar{a}_i = \bar{a}_i \bar{a}_1 \quad \text{if } i \neq 3$$

$$(D3) \quad \bar{a}_1 \bar{a}_3 \bar{a}_1 = \bar{a}_3 \bar{a}_1 \bar{a}_3$$

$$(D4) \quad \bar{a}_i \bar{a}_{i+1} \bar{a}_i = \bar{a}_{i+1} \bar{a}_i \bar{a}_{i+1} \quad \text{if } i = 2, 3, \dots, n - 1$$

$$(D5) \quad \bar{a}_i \bar{a}_j = \bar{a}_j \bar{a}_i \quad \text{if } |i - j| > 1$$

For $n \geq 4$, it is known that (D1)–(D5) are defining relations of the Hecke algebra $\mathcal{H}_{R,D_n}(q)$ of type D_n , so we may identify $\mathcal{H}_{R,D_n}(q)$ with $\bar{\mathcal{H}}_{R,B_n}(1, q)$.

Let u and q be indeterminates and

$$R_0 = \mathbb{Z}[u^{\pm 1}, q^{\pm 1}], \quad R_1 = \mathbb{Z}[q^{\pm 1}].$$

It is known that $\mathcal{H}_{R_0,B_n}(u, q)$ is a free R_0 -module of rank $2^n n!$ [1, 4] and that $\bar{\mathcal{H}}_{R_1,B_n}(1, q)$ is a free R_1 -module of rank $2^{n-1} n!$. Accordingly, $\mathcal{H}_{R_1,B_n}(1, q)$ is also a free R_1 -module of rank $2^n n!$.

4 Fixed subalgebras of $\mathcal{H}_{R_3,B_n}(1, q)$ by involutions

It is obvious that there is an algebra automorphism $\sharp$ of $\mathcal{H}_{R_1,B_n}(1, q)$ of order 2 which is defined by $a_1^\sharp = -a_1$ and $a_i^\sharp = a_i$ ($i \geq 2$). We define $\mathcal{H}_{R_1,B_n}(1, q)^\sharp$ to be the subalgebra consisting of fixed elements of $\sharp$ as follows.

$$\mathcal{H}_{R_1,B_n}(1, q)^\sharp = \{h \in \mathcal{H}_{R_1,B_n}(1, q) \mid h^\sharp = h\}.$$

The relations (B1') and (B2)–(B5) imply that every monomial with occurrences of even (resp. odd) numbers of a_1 also has occurrences of even (resp. odd) numbers of a_1 in each term of any other expression of it. Therefore $\mathcal{H}_{R_1, B_n}(1, q)^\sharp$ is generated by monomials with occurrences of even numbers of a_1 . Since the Weyl group W_{D_n} of type D_n is a normal subgroup of the Weyl group W_{B_n} of type B_n of index 2, and consists of all elements which can be written as products of $s_1, s_2, \dots, s_n$ with even numbers of factors of s_1 . On the other hand, W_{D_n} is generated by $s_1 s_2 s_1, s_2, s_3, \dots, s_n$ (see for example [3], 1.4.8). Thus we have the following direct sum of R_1 -modules:

$$\mathcal{H}_{R_1, B_n}(1, q)^\sharp = \bigoplus_{w \in W_{D_n}} R_1 T_w = \tilde{\mathcal{H}}_{R_1, B_n}(1, q).$$

The argument above is valid even if $n = 2, 3$. $\bigoplus_{w \in W_{D_n}} R T_w$ is closed under the product by (B1') and (B2)–(B5), Hence $\tilde{\mathcal{H}}_{R_1, B_n}(1, q) = \mathcal{H}_{R_1, B_n}(1, q)^\sharp$ holds. Let R_2, R_3 be the rings defined by

$$R_2 = \mathbb{Z}[u^{\pm 1}, q^{\pm 1}, (u+1)^{-1}, (q+1)^{-1}, \frac{1}{2}]$$

$$R_3 = \mathbb{Z}[q^{\pm 1}, (q+1)^{-1}, \frac{1}{2}].$$

It is known that there exists an algebra automorphism $\natural$ of $\mathcal{H}_{R_0, B_n}(u, q)$ of order 2 which is defined by $a_1^\natural = (u-1) - a_1$ and $a_i^\natural = (q-1) - a_i$ ($i \geq 2$). $\natural$ is called Goldman involution. We consider the fixed subalgebra by $\natural$ over R_2 . We define the elements b_i ($i = 1, 2, \dots, n$) of $\mathcal{H}_{R_2, B_n}(u, q)$ by

$$b_1 = \frac{a_1 - a_1^\natural}{u+1}, \quad b_i = \frac{a_i - a_i^\natural}{q+1} \quad \text{if } i = 2, 3, \dots, n$$

Then the following holds.

Proposition 4.1 ([6], Proposition 3.2). $b_1, b_2, \dots, b_n$ generate $\mathcal{H}_{R_2, B_n}(u, q)$ and constitute with the relations

$$(B'1) \quad b_i^2 = 1 \quad \text{if } i = 1, 2, \dots, n$$

$$(B'2) \quad b_1 b_2 b_1 b_2 = b_2 b_1 b_2 b_1 - 2 \frac{(u-1)(q-1)}{(u+1)(q+1)} (b_1 b_2 - b_2 b_1)$$

$$(B'3) \quad b_i b_{i+1} b_i = b_{i+1} b_i b_{i+1} - \left(\frac{q-1}{q+1} \right)^2 (b_i - b_{i+1}) \quad \text{if } i = 2, 3, \dots, n$$

$$(B'4) \quad b_i b_j = b_j b_i \quad \text{if } |i - j| > 1$$

a presentation of $\mathcal{H}_{R_2, B_n}(u, q)$.

Consider the following sets of monomials.

$$\begin{aligned}
\mathcal{S}_1 &= \{1, b_1\} \\
\mathcal{S}_2 &= \{1, b_2, b_2b_1, b_2b_1b_2\} \\
&\vdots \\
\mathcal{S}_i &= \{1, b_i, b_ib_{i-1}, \dots, b_ib_{i-1} \cdots b_2b_1, b_ib_{i-1} \cdots b_2b_1b_2, \dots, \\
&\quad b_ib_{i-1} \cdots b_2b_1b_2 \cdots b_{i-1}b_i\} \\
&\vdots \\
\mathcal{S}_n &= \{1, b_n, b_nb_{n-1}, \dots, b_nb_{n-1} \cdots b_2b_1, b_nb_{n-1} \cdots b_2b_1b_2, \dots, \\
&\quad b_nb_{n-1} \cdots b_2b_1b_2 \cdots b_{n-1}b_n\}
\end{aligned}$$

We shall say that $M = U_1U_2 \cdots U_n$ is a *monomial in b_i -normal form* in $\mathcal{H}_{R_2, B_n}(u, q)$, if $U_i \in \mathcal{S}_i$ for $i = 1, 2, \dots, n$. Then we have

Proposition 4.2 ([6], Proposition 3.3).

$$\mathcal{H}_{R_2, B_n}(u, q) = \bigoplus_{U_i \in \mathcal{S}_i} R_2U_1U_2 \cdots U_n$$

Applying the case $u = 1$ to the above proposition, we also obtain

$$\mathcal{H}_{R_3, B_n}(1, q) = \bigoplus_{U_i \in \mathcal{S}_i} R_3U_1U_2 \cdots U_n.$$

One can determine the fixed subalgebra by $\natural$ in $\mathcal{H}_{R_2, B_n}(u, q)$. Let $\mathcal{E}_n^\natural$ be the set of all monomials in b_i -normal form in $\mathcal{H}_{R_2, B_n}(u, q)$ which are products of even numbers of b_i 's. We define $\mathcal{H}_{R_2, B_n}(u, q)^\natural$ to be the R_2 -submodule of $\mathcal{H}_{R_2, B_n}(u, q)$ defined as follows.

$$\mathcal{H}_{R_2, B_n}(u, q)^\natural = \bigoplus_{M \in \mathcal{E}_n^\natural} R_2M$$

Proposition 4.3 ([6], Proposition 3.6). $\mathcal{H}_{R_2, B_n}(u, q)^\natural$ is the subalgebra of $\mathcal{H}_{R_2, B_n}(u, q)$ which consists of all the elements fixed under $\natural$. Furthermore $\text{rank}_{R_2} \mathcal{H}_{R_2, B_n}(u, q)^\natural = 2^{n-1}n!$.

We mention that this proposition may be applied to the case $u = 1$, and yields the same assertion for $\mathcal{H}_{R_3, B_n}(1, q)$ and $\mathcal{H}_{R_3, B_n}(1, q)^\natural$. In this case, $\natural$ turns out to be the automorphism determined by $a_1^\natural = -a_1$ and $a_i^\natural = (q-1) - a_i$ ($i \geq 2$). Let $\mathcal{S}_n$ be the set of all monomials in b_i -normal form in $\mathcal{H}_{R_2, B_n}(u, q)$. We define $\mathcal{H}_{R_2, B_n}(u, q)^{-\natural}$ to be the R_2 -submodule of $\mathcal{H}_{R_2, B_n}(u, q)$ as follows.

$$\mathcal{H}_{R_2, B_n}(u, q)^{-\natural} = \bigoplus_{M \in \mathcal{S}_n \setminus \mathcal{E}_n^\natural} R_2M$$

Then the following statement holds.

Proposition 4.4 ([6], Proposition 3.7). $\mathcal{H}_{R_2, B_n}(u, q)$ is a $\mathbb{Z}_2$ -crossed product with R_2 -submodules $A_{\bar{0}} = \mathcal{H}_{R_2, B_n}(u, q)^{\natural}$, $A_{\bar{1}} = \mathcal{H}_{R_2, B_n}(u, q)^{-\natural}$ and units $1 \in A_{\bar{0}}$, $b_1 \in A_{\bar{1}}$.

If we take $u = 1$, we obtain a $\mathbb{Z}_2$ -crossed product

$$\mathcal{H}_{R_3, B_n}(1, q) = \mathcal{H}_{R_3, B_n}(1, q)^{\natural} \oplus \mathcal{H}_{R_3, B_n}(1, q)^{-\natural}$$

Besides $\sharp, \natural$, we define one more algebra automorphism of $\mathcal{H}_{R_3, B_n}(1, q)$ of order 2. Let $\flat$ be the map defined by $a_1^{\flat} = a_1$ and $a_i^{\flat} = (q - 1) - a_i$ ($i \geq 2$). Then $\flat$ can be extended to an algebra automorphism of $\mathcal{H}_{R_3, B_n}(1, q)$. One can readily see that $\sharp \cdot \flat = \flat \cdot \sharp = \natural$. We consider the fixed subalgebra by $\flat$ over R_3 . We define $\mathcal{H}_{R_3, B_n}(1, q)^{\flat}$ to be the subalgebra consisting of fixed elements of $\flat$ as follows.

$$\mathcal{H}_{R_3, B_n}(1, q)^{\flat} = \{h \in \mathcal{H}_{R_3, B_n}(1, q) \mid h^{\flat} = h\}.$$

Let $\mathcal{E}_n^{\flat}$ be the set of all monomials in b_i -normal form in $\mathcal{H}_{R_3, B_n}(1, q)^{\flat}$ which have occurrences of even numbers of b_i ($i \neq 1$). For $n \geq 1$, we define $\tilde{\mathcal{H}}_{R_3, B_n}(1, q)$ to be the R_3 -submodule of $\mathcal{H}_{R_3, B_n}(1, q)$ defined as follows.

$$\tilde{\mathcal{H}}_{R_3, B_n}(1, q) = \bigoplus_{M \in \mathcal{E}_n^{\flat}} R_3 M$$

We have $b_1^{\flat} = b_1$ and $b_i^{\flat} = -b_i$ ($i \geq 2$) immediately. Hence the following holds.

Proposition 4.5. $\mathcal{H}_{R_3, B_n}(1, q)^{\flat} = \tilde{\mathcal{H}}_{R_3, B_n}(1, q)$. Furthermore, $\text{rank}_{R_3} \mathcal{H}_{R_3, B_n}(1, q)^{\flat} = 2^{n-1} n!$.

Proof. Equality of two algebras is by Proposition 4.1. We notice that $|\mathcal{S}_i| = 2i$. Let $\mathcal{S}_i^{\circ}$ be the subset of $\mathcal{S}_i$ consisting of the monomials with occurrences of odd numbers of b_i ($i \neq 1$). $|\mathcal{S}_i^{\circ}|$ is just i if $i > 1$ and $|\mathcal{S}_1^{\circ}| = 0$. A monomial $U_1 U_2 \cdots U_n$ in b_i -normal form has occurrences of even numbers of b_i ($i \neq 1$) if and only if the number of U_i such that $U_i \in \mathcal{S}_i^{\circ}$ is even. U_1 never belongs to $\mathcal{S}_1^{\circ}$. Hence there are 2^{n-2} cases of being so. In each case, there are $2n!$ ways of taking elements from $\mathcal{S}_i$'s. Thus, we have that there exist $2^{n-1} n!$ monomials in b_i -normal form with occurrences of even numbers of b_i ($i \neq 1$). $\square$

We define $\mathcal{H}_{R_3, B_n}(1, q)^{-\flat}$ to be the R_3 -submodule of $\mathcal{H}_{R_3, B_n}(1, q)$ as follows.

$$\mathcal{H}_{R_3, B_n}(1, q)^{-\flat} = \{h \in \mathcal{H}_{R_3, B_n}(1, q) \mid h^{\flat} = -h\}.$$

Then by Proposition 4.1, we have

$$\mathcal{H}_{R_3, B_n}(1, q)^{-\flat} = \bigoplus_{M \in \mathcal{S}_n \setminus \mathcal{E}_n^{\flat}} R_3 M.$$

Proposition 4.6. $\mathcal{H}_{R_3, B_n}(1, q)$ is a $\mathbb{Z}_2$ -crossed product with R_3 -submodules $A_{\bar{0}} = \mathcal{H}_{R_3, B_n}(1, q)^{\flat}$, $A_{\bar{1}} = \mathcal{H}_{R_3, B_n}(1, q)^{-\flat}$ and units $1 \in A_{\bar{0}}$, $b_2 \in A_{\bar{1}}$.

Proof. Clearly

$$A_{\bar{0}}A_{\bar{0}} = A_{\bar{0}}, \quad A_{\bar{1}}A_{\bar{1}} = A_{\bar{0}}, \quad A_{\bar{0}}A_{\bar{1}} = A_{\bar{1}}A_{\bar{0}} = A_{\bar{1}}$$

hold. Hence (C1) is satisfied. We may take units $1 \in A_{\bar{0}}$ and $b_2 \in A_{\bar{1}}$ so that $b_2A_{\bar{0}} = A_{\bar{0}}b_2 = A_{\bar{1}}$. Therefore (C2) and (C4) hold. Since $\mathcal{H}_{R_3, B_n}(1, q)^b = \bigoplus_{M \in \mathcal{E}_n^b} R_3M$ and $\mathcal{H}_{R_3, B_n}(1, q)^{-b} = \bigoplus_{M \in \mathcal{S}_n \setminus \mathcal{E}_n^b} R_3M$, (C3) holds. $\square$

Now we apply the generators b_i ($i = 1, \dots, n$) to $\mathcal{H}_{R_3, B_n}(1, q)^\sharp$. We immediately have $b_1 = a_1$ and $b_1^\sharp = -b_1$, $b_i^\sharp = b_i$ ($i = 2, \dots, n$). Let $\mathcal{E}_n^\sharp$ be the set of all monomials in b_i -normal form in $\mathcal{H}_{R_3, B_n}(1, q)$ which have occurrences of even numbers of b_1 . Then by Proposition 4.1, we immediately have

$$\begin{aligned} \mathcal{H}_{R_3, B_n}(1, q)^\sharp &= \bigoplus_{M \in \mathcal{E}_n^\sharp} R_3M \\ \mathcal{H}_{R_3, B_n}(1, q)^{-\sharp} &= \bigoplus_{M \in \mathcal{S}_n \setminus \mathcal{E}_n^\sharp} R_3M \end{aligned}$$

for $n \geq 1$. Moreover we obtain

Proposition 4.7. $\mathcal{H}_{R_3, B_n}(1, q)$ is a $\mathbb{Z}_2$ -crossed product with R_3 -submodules $A_{\bar{0}} = \mathcal{H}_{R_3, B_n}(1, q)^\sharp$, $A_{\bar{1}} = \mathcal{H}_{R_3, B_n}(1, q)^{-\sharp}$ and units $1 \in A_{\bar{0}}$, $b_1 \in A_{\bar{1}}$.

Proof. One can obtain the proof in the same manner as in Proposition 4.6. $\square$

5 A fixed subalgebra of $\mathcal{H}_{R_3, B_n}(1, q)^\sharp$ by Goldman involution

Let $\bar{b}_1 = b_1b_2b_1$, $\bar{b}_i = b_i$ ($i > 1$). By a direct computation, we get the following.

Proposition 5.1. $\bar{b}_1, \bar{b}_2, \dots, \bar{b}_n$ generate $\mathcal{H}_{R_3, B_n}(1, q)^\sharp$ and constitute with the relations

$$(D'1) \quad \bar{b}_i^2 = 1 \quad \text{if } i = 1, 2, \dots, n$$

$$(D'2) \quad \bar{b}_1\bar{b}_i = \bar{b}_i\bar{b}_1 \quad \text{if } i \neq 3$$

$$(D'3) \quad \bar{b}_1\bar{b}_3\bar{b}_1 = \bar{b}_3\bar{b}_1\bar{b}_3 - \left(\frac{q-1}{q+1}\right)^2(\bar{b}_1 - \bar{b}_3)$$

$$(D'4) \quad \bar{b}_i\bar{b}_{i+1}\bar{b}_i = \bar{b}_{i+1}\bar{b}_i\bar{b}_{i+1} - \left(\frac{q-1}{q+1}\right)^2(\bar{b}_i - \bar{b}_{i+1}) \quad \text{if } i = 2, 3, \dots, n-1$$

$$(D'5) \quad \bar{b}_i\bar{b}_j = \bar{b}_j\bar{b}_i \quad \text{if } |i-j| > 1$$

a presentation of $\mathcal{H}_{R_3, B_n}(1, q)^\sharp$.

The relations (D'1)–(D'5) imply that every product of an even (resp. odd) number of generators $\bar{b}_1, \bar{b}_2, \dots, \bar{b}_n$ must be written as a linear combination of products of even (resp. odd) numbers of the generators in any other expression of it. Therefore we define $\mathcal{H}_{R_3, B_n}(1, q)^\dagger$ to be the R_3 -subalgebra of $\mathcal{H}_{R_3, B_n}(1, q)^\sharp$ generated by all the monomials of even numbers of factors of $\bar{b}_1, \bar{b}_2, \dots, \bar{b}_n$. Goldman involution of $\mathcal{H}_{R_3, B_n}(1, q)^\sharp$ is given by $\bar{a}_i^\sharp = (q-1) - \bar{a}_i$ ($i = 1, 2, \dots, n$). We readily see that $\bar{b}_i^\sharp = -\bar{b}_i$. Thus the fixed subalgebra $(\mathcal{H}_{R_3, B_n}(1, q)^\sharp)^\natural$ of $\natural$ coincides with $\mathcal{H}_{R_3, B_n}(1, q)^\dagger$.

We consider intersections of these subalgebras $\mathcal{H}_{R_3, B_n}(1, q)^\natural$, $\mathcal{H}_{R_3, B_n}(1, q)^\flat$ and $\mathcal{H}_{R_3, B_n}(1, q)^\sharp$. We immediately have

$$\mathcal{H}_{R_3, B_n}(1, q)^\dagger = (\mathcal{H}_{R_3, B_n}(1, q)^\sharp)^\natural = \mathcal{H}_{R_3, B_n}(1, q)^\sharp \cap \mathcal{H}_{R_3, B_n}(1, q)^\natural.$$

Proposition 5.2.

$$\begin{aligned} \mathcal{H}_{R_3, B_n}(1, q)^\natural \cap \mathcal{H}_{R_3, B_n}(1, q)^\flat &= \mathcal{H}_{R_3, B_n}(1, q)^\flat \cap \mathcal{H}_{R_3, B_n}(1, q)^\sharp \\ &= \mathcal{H}_{R_3, B_n}(1, q)^\sharp \cap \mathcal{H}_{R_3, B_n}(1, q)^\natural \end{aligned}$$

Proof. Since $\flat = \natural \cdot \sharp$, we have

$$\begin{aligned} \mathcal{H}_{R_3, B_n}(1, q)^\natural \cap \mathcal{H}_{R_3, B_n}(1, q)^\flat &= \mathcal{H}_{R_3, B_n}(1, q)^\natural \cap \mathcal{H}_{R_3, B_n}(1, q)^{\natural \cdot \sharp} \\ &\subseteq \mathcal{H}_{R_3, B_n}(1, q)^\natural \cap \mathcal{H}_{R_3, B_n}(1, q)^\sharp \end{aligned}$$

Since $\sharp = \natural \cdot \flat$, we also have

$$\begin{aligned} \mathcal{H}_{R_3, B_n}(1, q)^\natural \cap \mathcal{H}_{R_3, B_n}(1, q)^\sharp &= \mathcal{H}_{R_3, B_n}(1, q)^\natural \cap \mathcal{H}_{R_3, B_n}(1, q)^{\natural \cdot \flat} \\ &\subseteq \mathcal{H}_{R_3, B_n}(1, q)^\natural \cap \mathcal{H}_{R_3, B_n}(1, q)^\flat \end{aligned}$$

Thus we obtain $\mathcal{H}_{R_3, B_n}(1, q)^\natural \cap \mathcal{H}_{R_3, B_n}(1, q)^\flat = \mathcal{H}_{R_3, B_n}(1, q)^\sharp \cap \mathcal{H}_{R_3, B_n}(1, q)^\natural$. Other equations are also proved in the same fashion. $\square$

From the relations in Proposition 4.1, we can see that every monomial with occurrences of even (resp. odd) numbers of a_1 also has occurrences of even (resp. odd) numbers of a_1 in each term of any other expression of it in $\mathcal{H}_{R_3, B_n}(1, q)$, and that the same is holds for numbers of occurrences of a_i 's ($i > 1$). Hence by the definition, $\mathcal{H}_{R_3, B_n}(1, q)^\dagger$ is generated by monomials with both occurrences of even numbers of a_1 and those of other a_i 's. We also have that

$$\mathcal{H}_{R_3, B_n}(1, q)^\dagger = \bigoplus_{M \in \mathcal{E}_n^\natural \cap \mathcal{E}_n^\natural} R_3 M.$$

If $n \geq 4$, then $\mathcal{H}_{R_3, B_n}(1, q)^\dagger$ yields to be the subalgebra $\mathcal{H}_{R_3, D_n}(q)^\natural$ of $\mathcal{H}_{R_3, D_n}(q)$ consisting of fixed elements of $\natural$.

Proposition 5.3. $\text{rank } \mathcal{H}_{R_3, B_n}(1, q)^\dagger = 2^{n-2}n!$.

Proof. We show this by induction on n . For $n = 2$, the assertion clearly holds. Assume $n > 2$. We define four subsets of $\mathcal{S}_n$ in $\mathcal{H}_{R_3, B_n}(1, q)$ as follows.

$$\begin{aligned} X_n &= \{M \mid M \text{ has an even number of } b_1 \text{ and an even number of } b_2, \dots, b_n\}, \\ Y_n &= \{M \mid M \text{ has an odd number of } b_1 \text{ and an even number of } b_2, \dots, b_n\}, \\ Z_n &= \{M \mid M \text{ has an even number of } b_1 \text{ and an odd number of } b_2, \dots, b_n\}, \\ W_n &= \{M \mid M \text{ has an odd number of } b_1 \text{ and an odd number of } b_2, \dots, b_n\}. \end{aligned}$$

Write $M = U_1 U_2 \cdots U_n$. We consider four cases depending upon the subset to which $M' = U_1 U_2 \cdots U_{n-1}$ belongs.

case 1 : If $M' \in X_{n-1}$, then U_n must have no b_1 and a even number of $b_2, \dots, b_n$.

case 2 : If $M' \in Y_{n-1}$, then U_n must have one b_1 and a even number of $b_2, \dots, b_n$.

case 3 : If $M' \in Z_{n-1}$, then U_n must have no b_1 and a odd number of $b_2, \dots, b_n$.

case 4 : If $M' \in W_{n-1}$, then U_n must have one b_1 and a odd number of $b_2, \dots, b_n$.

If n is even, exactly $n/2$ such U_n exist in each case. Thus we obtain

$$\begin{aligned} |X_n| &= \frac{n}{2}|X_{n-1}| + \frac{n}{2}|Y_{n-1}| + \frac{n}{2}|Z_{n-1}| + \frac{n}{2}|W_{n-1}| \\ &= \frac{n}{2}|\mathcal{S}_{n-1}| = \frac{n}{2}2^{n-1}(n-1)! = 2^{n-2}n!. \end{aligned}$$

If n is odd, then exactly $(n+1)/2$ such U_n exist in case 1,2 and $(n-1)/2$ in the other cases. Hence, we have

$$\begin{aligned} |X_n| &= \frac{n+1}{2}|X_{n-1}| + \frac{n+1}{2}|Y_{n-1}| + \frac{n-1}{2}|Z_{n-1}| + \frac{n-1}{2}|W_{n-1}| \\ &= \frac{n}{2}|\mathcal{S}_{n-1}| + \frac{1}{2}(|X_{n-1}| + |Y_{n-1}| - |Z_{n-1}| - |W_{n-1}|). \end{aligned}$$

Observing $|X_{n-1}| + |Y_{n-1}| = |\mathcal{E}_{n-1}^b| = 2^{n-2}(n-1)!$, we have $|Z_{n-1}| + |W_{n-1}| = 2^{n-1}(n-1)! - 2^{n-2}(n-1)! = 2^{n-2}(n-1)!$. Thus $|X_n| = 2^{-1}n|\mathcal{S}_{n-1}| = 2^{n-2}n!$ as desired. $\square$

Theorem 5.4. $\mathcal{H}_{R_3, B_n}(1, q)^\sharp$, $\mathcal{H}_{R_3, B_n}(1, q)^b$ and $\mathcal{H}_{R_3, B_n}(1, q)^\natural$ are $\mathbb{Z}_2$ -crossed products with $A_0 = \mathcal{H}_{R_3, B_n}(1, q)^\dagger$.

Proof. Let us define three submodules of $\mathcal{H}_{R_3, B_n}(1, q)$ as follows.

$$\begin{aligned} \mathcal{H}_{R_3, B_n}(1, q)^{-\dagger, \sharp} &= \bigoplus_{M \in Y_n \cup W_n} R_3 M, \\ \mathcal{H}_{R_3, B_n}(1, q)^{-\dagger, b} &= \bigoplus_{M \in Z_n \cup W_n} R_3 M, \\ \mathcal{H}_{R_3, B_n}(1, q)^{-\dagger, \natural} &= \bigoplus_{M \in Y_n \cup Z_n} R_3 M. \end{aligned}$$

It is clear that there exist direct sum decompositions as follows.

$$\begin{aligned}\mathcal{H}_{R_3, B_n}(1, q)^\sharp &= \mathcal{H}_{R_3, B_n}(1, q)^\dagger \oplus \mathcal{H}_{R_3, B_n}(1, q)^{-\dagger, \sharp}, \\ \mathcal{H}_{R_3, B_n}(1, q)^\flat &= \mathcal{H}_{R_3, B_n}(1, q)^\dagger \oplus \mathcal{H}_{R_3, B_n}(1, q)^{-\dagger, \flat}, \\ \mathcal{H}_{R_3, B_n}(1, q)^\natural &= \mathcal{H}_{R_3, B_n}(1, q)^\dagger \oplus \mathcal{H}_{R_3, B_n}(1, q)^{-\dagger, \natural}.\end{aligned}$$

One can readily check that these direct sum decompositions satisfy the condition of the $\mathbb{Z}_2$ -crossed product. $\square$

6 The branching rules for the Hecke algebras of Type B

Let $K_0 = \mathbb{Q}(u, q)$ be the quotient field of R_0 and $K_1 = \mathbb{Q}(q)$ that of R_1 . Let $\lambda = (\lambda^{(1)}, \lambda^{(2)})$ be a 2-tuple of Young diagrams of total size n . Let $T = (T^{(1)}, T^{(2)})$ be a tableau of shape λ . We mean by a tableau $T = (T^{(1)}, T^{(2)})$ of shape $\lambda = (\lambda^{(1)}, \lambda^{(2)})$ the shape of $T^{(i)}$ is $\lambda^{(i)}$ ($i = 1, 2$), and each of the symbols $1, 2, \dots, n$ appears in T exactly once. For $\lambda = (\lambda^{(1)}, \lambda^{(2)})$ (resp. $T = (T^{(1)}, T^{(2)})$), let $\lambda^{(i)'} (resp. T^{(i)'})$ be its transpose for $i = 1, 2$, and $\lambda' = (\lambda^{(1)'}, \lambda^{(2)'})$ (resp. $T' = (T^{(1)'}, T^{(2)'})$). Moreover, Let $\lambda^* = (\lambda^{(2)}, \lambda^{(1)})$, $T^* = (T^{(2)}, T^{(1)})$ and $\lambda'^* = (\lambda^{(2)'}, \lambda^{(1)'})$, $T'^* = (T^{(2)'}, T^{(1)'})$. A tableau of shape $\lambda = (\lambda^{(1)}, \lambda^{(2)})$ is said to be *standard* if the numbers $1, 2, \dots, n$ increase along the rows and columns of each Young diagram $\lambda^{(1)}, \lambda^{(2)}$ of λ . We denote by $T^{(\alpha)}(i, j)$ ($\alpha = 1, 2$) the number at i -th row and j -th column of $T^{(\alpha)}$. Let $\text{STab}(\lambda)$ be the set of all of standard tableaux of shape λ . Let k be the number at i -th row and j -th column of $T^{(1)}$ or $T^{(2)}$ of $T \in \text{STab}(\lambda)$, and l at i' -th row and j' -th column of them. Then the integer $d_{T, k, l} = (j' - i') - (j - i)$ is said to be *the axial distance from k to l in T* . For each $T \in \text{STab}(\lambda)$ we take a symbol v_T , and define the free K_0 -module

$$V_\lambda = \bigoplus_{T \in \text{STab}(\lambda)} K_0 v_T.$$

For each λ , we can give an irreducible representation of $\mathcal{H}_{K_0, B_n}(u, q)$ in the following manner. Let y be any indeterminate and k any integer. Let $M(k, y)$ be the 2×2 matrix defined by

$$M(k, y) = \frac{1}{1 - q^k y} \begin{bmatrix} q - 1 & 1 - q^{k+1} y \\ q(1 - q^{k-1} y) & -q^k y(q - 1) \end{bmatrix}$$

For a standard tableau $T = (T^{(1)}, T^{(2)})$ of shape $\lambda = (\lambda^{(1)}, \lambda^{(2)})$ of total size n , we define ρ_T to be the map from $\{1, 2, \dots, n\}$ to $\{1, 2\}$ such that the number i occurs in the $\rho_T(i)$ -th Young diagram $T^{(\rho_T(i))}$ of T . We set $u_1 = u$ and $u_2 = -1$.

We shall give an action of the generators of $\mathcal{H}_{K_0, B_n}(u, q)$ on V_λ as follows.

- (a1) $a_1 v_T = u_{\rho_T(1)} v_T$.
- (a2) If $i > 1$, a_i acts on V_λ in three ways, depending upon the position that $i - 1$ and i occupy in $T = (T^{(1)}, T^{(2)})$ as follows.

- (a) If $i - 1$ and i appear in the same row of the same diagram of $(T^{(1)}, T^{(2)})$, then $a_i v_T = q v_T$.
- (b) If $i - 1$ and i appear in the same column of the same diagram of $(T^{(1)}, T^{(2)})$, then $a_i v_T = -v_T$.
- (c) Elsewhere, a_i acts on the subspace $K_0 v_T \oplus K_0 v_{s_{i-1}T}$ of V_λ as follows, where $s_{i-1} = (i - 1, i)$ is the transposition of $i - 1$ and i , and $s_{i-1}T$ is the standard tableau obtained from T by transposing $i - 1$ and i .

$$a_i \langle v_T, v_{s_{i-1}T} \rangle = \langle v_T, v_{s_{i-1}T} \rangle M \left(d_{T,i,i-1}, \frac{u_{\rho_T(i-1)}}{u_{\rho_T(i)}} \right).$$

Theorem 6.1 ([4], Theorem 2.2.11). *For each 2-tuple of Young diagrams of total size n , the above action of $\mathcal{H}_{K_0, B_n}(u, q)$ gives an (absolutely) irreducible representation of $\mathcal{H}_{K_0, B_n}(u, q)$. If $\lambda \neq \mu$ as 2-tuples of Young diagrams, then V_λ and V_μ are mutually inequivalent irreducible representations of $\mathcal{H}_{K_0, B_n}(u, q)$. These constitute a complete set of representatives of irreducible representations of $\mathcal{H}_{K_0, B_n}(u, q)$*

These representations are said to be the *seminormal form representations* of $\mathcal{H}_{K_0, B_n}(u, q)$. We denote by (π_λ, V_λ) the seminormal form representation of $\mathcal{H}_{K_0, B_n}(u, q)$ corresponding to λ . By an easy calculation, we obtain that $\pi_\lambda(b_i)$ is as follows.

- (b1) $\pi_\lambda(b_1) v_T = (-1)^{\rho_T(1)-1} v_T$.
- (b2) If $i > 1$, $\pi_\lambda(b_i)$ are given in three ways, depending upon the position that $i - 1$ and i occupy in $T = (T^{(1)}, T^{(2)})$ as follows.
 - (a) If $i - 1$ and i appear in the same row of the same diagram of $(T^{(1)}, T^{(2)})$, then $\pi_\lambda(b_i) v_T = v_T$.
 - (b) If $i - 1$ and i appear in the same column of the same diagram of $(T^{(1)}, T^{(2)})$, then $\pi_\lambda(b_i) v_T = -v_T$.
 - (c) Elsewhere, $\pi_\lambda(b_i)$ acts on the subspace $K_0 v_T \oplus K_0 v_{s_{i-1}T}$ of V_λ as follows.

$$\pi_\lambda(b_i) \langle v_T, v_{s_{i-1}T} \rangle = \langle v_T, v_{s_{i-1}T} \rangle M' \left(d_{T,i,i-1}, \frac{u_{\rho_T(i-1)}}{u_{\rho_T(i)}} \right)$$

where $M'(k, y)$ is a 2×2 matrix defined by

$$M'(k, y) = \frac{1}{(q+1)(1-q^k y)} \begin{bmatrix} (q-1)(1+q^k y) & 2(1-q^{k+1} y) \\ 2q(1-q^{k-1} y) & -(q-1)(1+q^k y) \end{bmatrix}$$

We denote by $(\pi_{\lambda, \mathfrak{h}}, V_{\lambda, \mathfrak{h}})$ the restriction of (π_λ, V_λ) to $\mathcal{H}_{K_0, B_n}(u, q)^{\mathfrak{h}}$. We obtained the branching rule between $\mathcal{H}_{K_0, B_n}(u, q)$ and $\mathcal{H}_{K_0, B_n}(u, q)^{\mathfrak{h}}$ in [6] as follows.

Theorem 6.2 ([6] Corollary 4.5, Proposition 4.7, Theorem 4.8). (1) $\pi_{\lambda'^*, \mathfrak{h}} \cong \pi_{\lambda, \mathfrak{h}}$.

- (2) If $\lambda' = \lambda$, then $\pi_{\lambda, \mathfrak{h}}$ decomposes into inequivalent subrepresentations $\pi_{\lambda, \mathfrak{h}}^+$ and $\pi_{\lambda, \mathfrak{h}}^-$ over $\bar{K}_0$, the algebraic closure of K_0 . Furthermore, $\deg \pi_{\lambda, \mathfrak{h}}^+ = \deg \pi_{\lambda, \mathfrak{h}}^- = \deg \pi_{\lambda, \mathfrak{h}}/2$.

(3) Let $\{\lambda_i, \lambda_i'^*, \mu_j\}_{i,j}$ be the set of 2-tuples of Young diagrams such that

$$\lambda_i'^* \neq \lambda_i, \quad \mu_j'^* = \mu_j$$

Then

$$\text{Irr}(\mathcal{H}_{\bar{K}_0, B_n}(q)^{\natural}) = \{\pi_{\lambda_i, \natural}, \pi_{\mu_j, \natural}^+, \pi_{\mu_j, \natural}^-\}_{i,j}$$

is a basic set of irreducible representations of $\mathcal{H}_{\bar{K}_0, B_n}(u, q)^{\natural}$. $\mathcal{H}_{\bar{K}_0, B_n}(u, q)^{\natural}$ is semisimple.

The proof is in [6], but there are some errors in it. So we shall prove again. From (b1),(b2), we obtain that $\pi_{\lambda'^*}(b_i)$ is as follows by a direct computation.

$$(c1) \quad \pi_{\lambda'^*}(b_1)v_{T'^*} = (-1)^{\rho_T(1)}v_{T'^*}.$$

(c2) If $i > 1$, $\pi_{\lambda'^*}(b_i)$ are given in three ways, depending upon the position that $i-1$ and i occupy in $T = (T^{(1)}, T^{(2)})$ as follows.

- (a) If $i-1$ and i appear in the same row of the same diagram of T , then $\pi_{\lambda'^*}(b_i)v_{T'^*} = -v_{T'^*}$.
- (b) If $i-1$ and i appear in the same column of the same diagram of T , then $\pi_{\lambda'^*}(b_i)v_{T'^*} = v_{T'^*}$.
- (c) Elsewhere, $\pi_{\lambda'^*}(b_i)$ reduces to the endomorphism of the subspace $K_0v_{T'^*} \oplus K_0v_{s_{i-1}T'^*}$ of $V_{\lambda'^*}$ as follows.

$$\begin{aligned} \pi_{\lambda'^*}(b_i)\langle v_{T'^*}, v_{s_{i-1}T'^*} \rangle &= \langle v_{T'^*}, v_{s_{i-1}T'^*} \rangle M' \left(-d_{T,i,i-1}, \frac{u_{\rho_T(i)}}{u_{\rho_T(i-1)}} \right) \\ &= \langle v_{T'^*}, v_{s_{i-1}T'^*} \rangle JM' \left(d_{T,i,i-1}, \frac{u_{\rho_T(i-1)}}{u_{\rho_T(i)}} \right) J \end{aligned}$$

$$\text{where } J = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

In [6], p.243-244, the intertwining operator between $\pi_{\lambda'^*, \natural}$ and $\pi_{\lambda, \natural}$ is given. But it is incorrect since the denominator of $\alpha_T(i, j)$ equals 0 if i and j belong to the same Young diagram and satisfy $d_{T,i,j} = 0$. So we correct the intertwining operator as follows. For each standard tableau T of shape λ of total size n , we define the map ψ_T from $I = \{(i, j) \in \mathbb{N} \times \mathbb{N} \mid 1 \leq i \neq j \leq n\}$ to K_0 divided into two cases as follows.

case 1 : i and j appear in the same Young diagram.

$$\psi_T(i, j) = \begin{cases} 1 & \text{if } i \text{ and } j \text{ appear in the same row or the same column,} \\ & \text{or } i \text{ and } j \text{ satisfy } d_{T,i,j} = 0, \\ \frac{q(1 - q^{d_{T,i,j}-1}u_{\rho_T(j)}u_{\rho_T(i)}^{-1})}{(q+1)(1 - q^{d_{T,i,j}}u_{\rho_T(j)}u_{\rho_T(i)}^{-1})} \times \text{sgn}(d_{T,i,j}) & \text{otherwise.} \end{cases}$$

case 2 : i and j appear in the different Young diagrams.

$$\psi_T(i, j) = \frac{q(1 - q^{d_{T,i,j}-1}u_{\rho_T(j)}u_{\rho_T(i)}^{-1})}{(q+1)(1 - q^{d_{T,i,j}}u_{\rho_T(j)}u_{\rho_T(i)}^{-1})} \times (\rho_T(i) - \rho_T(j))$$

Let $\psi : \text{STab}(\lambda) \longrightarrow K_0$ be the map defined by

$$\psi(T) = \prod_{1 \leq j < i \leq n} \psi_T(i, j), \quad (1)$$

and $\Psi_{\natural}$ the K_0 -homomorphism defined by

$$\begin{aligned} \Psi_{\natural} : V_{\lambda} &\longrightarrow V_{\lambda'^*} \\ v_T &\longmapsto \psi(T)v_{T'^*} \end{aligned}$$

Then, Proposition 4.4 in [6] holds correctly as follows.

Proposition 6.3.

$$\pi_{\lambda'^*}(b_i)\Psi_{\natural}(v_T) = - \sum_{S \in \text{STab}(\lambda)} \left(\pi_{\lambda}(b_i) \right)_{T,S} \Psi_{\natural}(v_S) \quad \text{for } i = 1, 2, \dots, n$$

Proof. In the cases (c1) and (c2)-(a)(b), the equation holds obviously. Hence we consider only the case (c2)-(c). By a direct calculation, we have

$$\begin{aligned} & \frac{(1 - q^{d_{T,i,i-1}+1} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})}{(q+1)(1 - q^{d_{T,i,i-1}} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})} \\ &= \frac{q(1 - q^{d_{s_{i-1}T,i,i-1}-1} u_{\rho_{s_{i-1}T}(i-1)} u_{\rho_{s_{i-1}T}(i)}^{-1})}{(q+1)(1 - q^{d_{s_{i-1}T,i,i-1}} u_{\rho_{s_{i-1}T}(i-1)} u_{\rho_{s_{i-1}T}(i)}^{-1})} \end{aligned}$$

Hence by the definition of $\psi(T)$, one can check the following.

$$\begin{aligned} & \frac{(1 - q^{d_{T,i,i-1}+1} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})}{(q+1)(1 - q^{d_{T,i,i-1}} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})} \psi(T) \\ &= - \frac{q(1 - q^{d_{T,i,i-1}-1} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})}{(q+1)(1 - q^{d_{T,i,i-1}} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})} \psi(s_{i-1}T) \end{aligned}$$

Therefore, we obtain the following equation.

$$\begin{aligned} \pi_{\lambda'^*}(b_i)\psi(T)v_{T'^*} &= \frac{(q-1)(1 + q^{d_{T'^*,i,i-1}} u_{\rho_{T'^*}(i-1)} u_{\rho_{T'^*}(i)}^{-1})}{(q+1)(1 - q^{d_{T'^*,i,i-1}} u_{\rho_{T'^*}(i-1)} u_{\rho_{T'^*}(i)}^{-1})} \psi(T)v_{T'^*} \\ &+ \frac{2q(1 - q^{d_{T'^*,i,i-1}-1} u_{\rho_{T'^*}(i-1)} u_{\rho_{T'^*}(i)}^{-1})}{(q+1)(1 - q^{d_{T'^*,i,i-1}} u_{\rho_{T'^*}(i-1)} u_{\rho_{T'^*}(i)}^{-1})} \psi(T)v_{s_{i-1}T'^*} \\ &= - \frac{(q-1)(1 + q^{d_{T,i,i-1}} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})}{(q+1)(1 - q^{d_{T,i,i-1}} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})} \psi(T)v_{T'^*} \\ &- \frac{2q(1 - q^{d_{T,i,i-1}-1} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})}{(q+1)(1 - q^{d_{T,i,i-1}} u_{\rho_T(i-1)} u_{\rho_T(i)}^{-1})} \psi(s_{i-1}T)v_{s_{i-1}T'^*} \end{aligned}$$

Thus we have proved the proposition. $\square$

Corollary 6.4. $\pi_{\lambda, \natural} \cong \pi_{\lambda'^*, \natural}$. The K_0 -homomorphism $\Psi_{\natural} : V_{\lambda, \natural} \longrightarrow V_{\lambda'^*, \natural}$ is the intertwining operator between $\pi_{\lambda, \natural}$ and $\pi_{\lambda'^*, \natural}$.

Proof. $\mathcal{H}_{K_0, B_n}(u, q)^{\natural}$ is generated by all the monomials of even numbers of products of $b_1, b_2, \dots, b_n$. Thus the representation matrix of $\pi_{\lambda, \natural}$ with respect to the basis $\{v_T | T \in \text{STab}(\lambda)\}$ and that of $\pi_{\lambda'^*, \natural}$ with respect to the basis $\{\Psi_{\natural}(v_T) | T \in \text{STab}(\lambda)\}$ coincide. $\square$

Therefore we obtain (1) of Theorem 6.2.

Corollary 6.5. Let $b \in \mathcal{H}_{K_0, B_n}(u, q)^{\natural}$. Then

$$\left(\pi_{\lambda'^*}(b) \right)_{T'^*, S'^*} \psi(T) = \left(\pi_{\lambda}(b) \right)_{T, S} \psi(S)$$

Proof. Let $b = b_{j_1} b_{j_2} \cdots b_{j_k} \in \mathcal{H}_{K_0, B_n}(u, q)^{\natural}$. Let $T_1 = T, T_k = S$. Then we may write

$$\pi_{\lambda'^*}(b) \psi(T_1) v_{T_1'^*} = \sum_{T_k \in \text{STab}(\lambda)} \left(\pi_{\lambda'^*}(b) \right)_{T_1'^*, T_k'^*} \psi(T_1) v_{T_k'^*}.$$

Since k is even, we obtain the following by using Proposition 6.3 repeatedly.

$$\begin{aligned} \pi_{\lambda'^*}(b) \psi(T_1) v_{T_1'^*} &= \pi_{\lambda'^*}(b_{j_1}) \pi_{\lambda'^*}(b_{j_2}) \cdots \pi_{\lambda'^*}(b_{j_k}) \psi(T_1) v_{T_1'^*} \\ &= \sum_{T_2, T_3, \dots, T_k \in \text{STab}(\lambda)} \left(\pi_{\lambda}(b_{j_k}) \right)_{T_1, T_2} \left(\pi_{\lambda}(b_{j_{k-1}}) \right)_{T_2, T_3} \cdots \left(\pi_{\lambda}(b_{j_1}) \right)_{T_{k-1}, T_k} \psi(T_k) v_{T_k'^*} \\ &= \sum_{T_k \in \text{STab}(\lambda)} \left(\pi_{\lambda}(b) \right)_{T_1, T_k} \psi(T_k) v_{T_k'^*}. \end{aligned}$$

Comparing the coefficients of $v_{T_k'^*}$ of the both right-hand sides, we obtain the assertion. $\square$

For λ such that $\lambda'^* = \lambda$, we define $\text{STab}(\lambda)^{\natural}$ and $\text{STab}(\lambda)^{-\natural}$ to be the sets of all the standard tableaux of shape λ such that 1 belongs to $T^{(1)}$ and $T^{(2)}$ respectively. Then, $\text{STab}(\lambda)$ is the disjoint union of $\text{STab}(\lambda)^{\natural}$ and $\text{STab}(\lambda)^{-\natural}$. There exists an involutive bijection from $\text{STab}(\lambda)^{\natural}$ to $\text{STab}(\lambda)^{-\natural}$ which is defined by $T \mapsto T'^*$. The submodules $V_{\lambda, \natural}^+, V_{\lambda, \natural}^-$ of $V_{\lambda, \natural}$ have also been given in p.246 of [6] as follows.

$$\begin{aligned} V_{\lambda, \natural}^+ &= \bigoplus_{T \in \text{STab}(\lambda)^{\natural}} \bar{K}_1(\sqrt{\psi(T'^*)} v_T + \sqrt{\psi(T)} v_{T'^*}), \\ V_{\lambda, \natural}^- &= \bigoplus_{T \in \text{STab}(\lambda)^{\natural}} \bar{K}_1(\sqrt{\psi(T'^*)} v_T - \sqrt{\psi(T)} v_{T'^*}), \end{aligned}$$

We notice that $\sqrt{\psi(T)}$ has two branches. We take a suitable branch in each computation of square roots for consistency. As in the proof of Proposition 4.7 of [6], we also have

$$\begin{aligned} \pi_{\lambda, \natural}(b) \left(\sqrt{\psi(T'^*)} v_T \pm \sqrt{\psi(T)} v_{T'^*} \right) &= \sum_{S \in \text{STab}(\lambda)^{\natural}} \left\{ \sqrt{\left(\pi_{\lambda, \natural}(b) \right)_{T, S} \left(\pi_{\lambda, \natural}(b) \right)_{T'^*, S'^*}} \right. \\ &\quad \left. \pm \sqrt{\left(\pi_{\lambda, \natural}(b) \right)_{T, S'^*} \left(\pi_{\lambda, \natural}(b) \right)_{T'^*, S}} \right\} \times \left(\sqrt{\psi(S'^*)} v_S + \sqrt{\psi(S)} v_{S'^*} \right) \end{aligned}$$

for $T \in \text{STab}(\lambda)^{\natural}$.

proof of Theorem 6.2. (1) is Corollary 6.4 itself. If $\lambda'^* = \lambda$, then one can deduce directly from Theorem 2.7 that both $\pi_{\lambda, \natural}^+$ and $\pi_{\lambda, \natural}^-$ are irreducible and mutually inequivalent. Thus (2) holds. Theorem 2.7 also shows immediately that $\pi_{\lambda_i, \natural}, \pi_{\mu_j, \natural}^+, \pi_{\mu_j, \natural}^-$ are mutually inequivalent. Semisimplicity of $\mathcal{H}_{\tilde{K}_0, B_n}(u, q)^\natural$ has been proved in Theorem 4.8 of [6]. $\square$

On the other hand, The branching rule between $\mathcal{H}_{K_1, B_n}(1, q)$ and $\mathcal{H}_{K_1, B_n}(1, q)^\natural$ has given in [4] for $n \geq 4$. We also refer [3] for the detail of the irreducible representations of $\mathcal{H}_{K_1, D_n}(q)$. We shall give a proof of the branching rule for $n \geq 2$ using the theory of crossed products. Now we set $u = 1$. We denote by $(\pi_{\lambda, \#}, V_{\lambda, \#})$ the restriction of (π_λ, V_λ) to $\mathcal{H}_{K_1, B_n}(1, q)^\natural$. From (b1), (b2), we obtain that $\pi_{\lambda^*}(b_i)$ is as follows.

$$(d1) \quad \pi_{\lambda^*}(b_1)v_{T^*} = (-1)^{\rho_T(1)}v_{T^*}.$$

(d2) If $i > 1$, $\pi_{\lambda^*}(b_i)$ are given in three ways, depending upon the position that $i-1$ and i occupy in $T = (T^{(1)}, T^{(2)})$ as follows.

- (a) If $i-1$ and i appear in the same row of the same diagram of T , then $\pi_{\lambda^*}(b_i)v_{T^*} = v_{T^*}$.
- (b) If $i-1$ and i appear in the same column of the same diagram of T , then $\pi_{\lambda^*}(b_i)v_{T^*} = -v_{T^*}$.
- (c) Elsewhere, $\pi_{\lambda^*}(b_i)$ acts on the subspace $K_0v_{T^*} \oplus K_0v_{s_{i-1}T^*}$ of V_{λ^*} as follows.

$$\pi_{\lambda^*}(b_i)\langle v_{T^*}, v_{s_{i-1}T^*} \rangle = \langle v_{T^*}, v_{s_{i-1}T^*} \rangle M' \left(d_{T, i, i-1}, \frac{u_{\rho_T(i-1)}}{u_{\rho_T(i)}} \right)$$

Indeed, (d1) and (d2)-(a)(b) are obvious, and (d2)-(c) follows from $d_{T^*, i, i-1} = d_{T, i, i-1}$, $u_{\rho_{T^*}(i-1)}/u_{\rho_{T^*}(i)} = u_{\rho_T(i-1)}/u_{\rho_T(i)}$. Therefore, (d1), (d2) give the following.

Proposition 6.6 ([3] 10.4). $\pi_{\lambda, \#} \cong \pi_{\lambda^*, \#}$. Especially, two matrices correspond to $\pi_{\lambda^*, \#}(h)$ and $\pi_{\lambda, \#}(h)$ ($h \in \mathcal{H}_{K_1, B_n}(1, q)^\natural$) coincide.

Proof. Since $\mathcal{H}_{K_1, B_n}(1, q)^\natural$ is generated by all the monomials with occurrences of even numbers of b_1 , all the matrices $\pi_{\lambda^*, \#}(h)$ ($h \in \mathcal{H}_{K_1, B_n}(1, q)^\natural$) coincide with $\pi_{\lambda, \#}(h)$. $\square$

If $\lambda^* = \lambda$, we shall show that $V_{\lambda, \#}$ decomposes into two nonzero submodules. Let ω be the endomorphism of V_λ defined by

$$\omega(v_T) = v_{T^*}$$

Then, obviously $\omega^2 = 1$. Furthermore, ω satisfies the following property.

Proposition 6.7. $\omega\pi_\lambda(h) = \pi_\lambda(h)\omega$ for all $h \in \mathcal{H}_{K_1, B_n}(1, q)^\natural$.

Proof. If $h = b_1$, then

$$\begin{aligned} \omega(\pi_\lambda(b_1)v_T) &= (-1)^{\rho_T(1)-1}\omega(v_T) = (-1)^{\rho_T(1)-1}v_{T^*} = -(-1)^{\rho_T(1)}v_{T^*} \\ \pi_\lambda(b_1)\omega(v_T) &= \pi_\lambda(b_1)v_{T^*} = (-1)^{\rho_T(1)}v_{T^*} \end{aligned}$$

Hence $\omega(\pi_\lambda(b_1)v_T) = -\pi_\lambda(b_1)\omega(v_T)$.

If $h = b_i$ ($i > 1$), then by (b2) and (d2) we have

$$\omega(\pi_\lambda(b_i)v_T) = \pi_\lambda(b_i)\omega(v_T).$$

Since $\mathcal{H}_{K_1, B_n}(1, q)^\sharp$ is generated by all the monomials with occurrences of even numbers of b_1 , $\omega\pi_\lambda(h) = \pi_\lambda(h)\omega$ holds. $\square$

Let $\lambda^* = \lambda$. We define $\text{STab}(\lambda)^\sharp$ and $\text{STab}(\lambda)^{-\sharp}$ to be the sets of all the standard tableaux T of shape λ such that 1 belongs to $T^{(1)}$ and $T^{(2)}$ respectively. Clearly $\text{STab}(\lambda)$ is the disjoint union of $\text{STab}(\lambda)^\sharp$ and $\text{STab}(\lambda)^{-\sharp}$. There exists an involutive bijection from $\text{STab}(\lambda)^\sharp$ to $\text{STab}(\lambda)^{-\sharp}$ which is defined by $T \mapsto T^*$. We define two subspaces $V_{\lambda, \sharp}^+, V_{\lambda, \sharp}^-$ as follows.

$$\begin{aligned} V_{\lambda, \sharp}^+ &= \bigoplus_{T \in \text{STab}(\lambda)^\sharp} K_1(v_T + v_{T^*}) \\ V_{\lambda, \sharp}^- &= \bigoplus_{T \in \text{STab}(\lambda)^\sharp} K_1(v_T - v_{T^*}) \end{aligned}$$

Then $\dim V_{\lambda, \sharp}^+ = \dim V_{\lambda, \sharp}^- = \dim V_\lambda/2$ and $V_\lambda = V_{\lambda, \sharp}^+ \oplus V_{\lambda, \sharp}^-$. Moreover, $V_{\lambda, \sharp}^+$ and $V_{\lambda, \sharp}^-$ are the eigenspaces corresponding to the eigenvalues 1 and -1 of ω respectively. Therefore, $V_{\lambda, \sharp}^+$ and $V_{\lambda, \sharp}^-$ are $\mathcal{H}_{K_1, B_n}(1, q)^\sharp$ -submodules. We denote by $(\pi_{\lambda, \sharp}^+, V_{\lambda, \sharp}^+)$ and $(\pi_{\lambda, \sharp}^-, V_{\lambda, \sharp}^-)$ the representations corresponding to the submodules $V_{\lambda, \sharp}^+, V_{\lambda, \sharp}^-$ respectively.

Theorem 6.8. (1) $\pi_{\lambda^*, \sharp} \cong \pi_{\lambda, \sharp}$.

(2) If $\lambda^* = \lambda$, then $\pi_{\lambda, \sharp}$ decomposes into inequivalent subrepresentations $\pi_{\lambda, \sharp}^+$ and $\pi_{\lambda, \sharp}^-$ over $\bar{K}_1$, the algebraic closure of K_1 . Furthermore, $\deg \pi_{\lambda, \sharp}^+ = \deg \pi_{\lambda, \sharp}^- = \deg \pi_{\lambda, \sharp}/2$.

(3) Let $\{\lambda_i, \lambda_i^*, \mu_j\}_{i,j}$ be the set of 2-tuples of Young diagrams such that

$$\lambda_i^* \neq \lambda_i, \quad \mu_j^* = \mu_j$$

Then

$$\text{Irr}(\mathcal{H}_{\bar{K}_1, B_n}(q)^\sharp) = \{\pi_{\lambda_i, \sharp}, \pi_{\mu_j, \sharp}^+, \pi_{\mu_j, \sharp}^-, \}_{i,j}$$

is a basic set of irreducible representations of $\mathcal{H}_{\bar{K}_1, B_n}(q)^\sharp$. $\mathcal{H}_{\bar{K}_1, B_n}(q)^\sharp$ is semisimple.

Proof. (1) is Proposition 6.6 itself. The submodules $V_{\lambda, \sharp}^+$ and $V_{\lambda, \sharp}^-$ correspond to the case (2) of Theorem 2.7. Hence they are simple and mutually non-isomorphic. Thus (2) holds. The proof of (3) is in the same manner as in the proof of Theorem 6.2, just replacing $\natural$ and $\bar{K}_0$ with $\sharp$ and $\bar{K}_1$ respectively. So we omit the detail. $\square$

Next we shall give the branching rule between $\mathcal{H}_{K_1, B_n}(1, q)$ and $\mathcal{H}_{K_1, B_n}(1, q)^\flat$. From (b1), (b2), we obtain that $\pi_{\lambda'}(b_i)$ is as follows.

(e1) $\pi_{\lambda'}(b_1)v_{T'} = (-1)^{\rho_T(1)-1}v_{T'}$.

(e2) If $i > 1$, $\pi_{\lambda'}(b_i)$ are given in three ways, depending upon the position that $i-1$ and i occupy in $T = (T^{(1)}, T^{(2)})$ as follows.

- (a) If $i-1$ and i appear in the same row of the same diagram of T , then $\pi_{\lambda'}(b_i)v_{T'} = -v_{T'}$.
- (b) If $i-1$ and i appear in the same column of the same diagram of T , then $\pi_{\lambda'}(b_i)v_{T'} = v_{T'}$.
- (c) Elsewhere, $\pi_{\lambda'}(b_i)$ acts on the subspace $K_0v_{T'} \oplus K_1v_{s_{i-1}T'}$ of $V_{\lambda'}$ as follows.

$$\pi_{\lambda'}(b_i)\langle v_{T'}, v_{s_{i-1}T'} \rangle = \langle v_{T'}, v_{s_{i-1}T'} \rangle M' \left(-d_{T,i,i-1}, \frac{u_{\rho_T(i-1)}}{u_{\rho_T(i)}} \right)$$

Indeed, (e1) is obvious, and the equations $d_{T',i,i-1} = -d_{T,i,i-1}$, $u_{\rho_{T'}(i-1)} = u_{\rho_T(i-1)}$, $u_{\rho_{T'}(i)} = u_{\rho_T(i)}$ imply (e2)-(c). We notice that $u_{\rho_T(i-1)}/u_{\rho_T(i)}$ equals 1 if i and $i-1$ belong to same Young diagram and -1 if not.

We shall give the intertwining operator between $\pi_{\lambda,b}$ and $\pi_{\lambda',b}$. Using $\psi(T)$ which has given in the equation (1), we define the K_1 -homomorphism Ψ_b to be

$$\begin{aligned} \Psi_b : V_{\lambda} &\longrightarrow V_{\lambda'} \\ v_T &\longmapsto \psi(T)v_{T'} \end{aligned}$$

We denote the action of $b \in \mathcal{H}_{K_1, B_n}(1, q)$ by

$$\pi_{\lambda}(b)v_T = \sum_{S \in \text{STab}(\lambda)} \left(\pi_{\lambda}(b) \right)_{T,S} v_S$$

where $(\pi_{\lambda}(b))_{T,S}$'s are elements in K_1 .

Proposition 6.9.

$$\begin{aligned} \pi_{\lambda'}(b_1)\Psi_b(v_T) &= \sum_{S \in \text{STab}(\lambda)} \left(\pi_{\lambda}(b_1) \right)_{T,S} \Psi_b(v_S) \\ \pi_{\lambda'}(b_i)\Psi_b(v_T) &= - \sum_{S \in \text{STab}(\lambda)} \left(\pi_{\lambda}(b_i) \right)_{T,S} \Psi_b(v_S) \quad \text{for } i = 2, \dots, n \end{aligned}$$

Proof. From (b1) and (e1), it is trivial for b_1 . Assume $i > 1$. Comparing (b2) and (e2), the assertion holds for the case (a),(b). For the case (c), observing that $u_{\rho_T(i-1)}/u_{\rho_T(i)}$ equals 1 or -1 , one can prove in the same manner as in the proof of Proposition 6.3, just replacing $'^*$ with $'$, so we omit the detail. $\square$

We denote by $(\pi_{\lambda,b}, V_{\lambda,b})$ the restriction of $(\pi_{\lambda}, V_{\lambda})$ to $\mathcal{H}_{K_1, B_n}(1, q)^b$. In the same fashions as in Corollary 6.4 and Corollary 6.5, we also obtain the following two corollaries.

Corollary 6.10. $\pi_{\lambda,b} \cong \pi_{\lambda',b}$. The K_1 -homomorphism $\Psi_b : V_{\lambda,b} \longrightarrow V_{\lambda',b}$ is the intertwining operator between $\pi_{\lambda,b}$ and $\pi_{\lambda',b}$.

Corollary 6.11. *Let $b \in \mathcal{H}_{K_1, B_n}(1, q)^b$. Then*

$$\left(\pi_{\lambda'}(b) \right)_{T', S'} \psi(T) = \left(\pi_{\lambda}(b) \right)_{T, S} \psi(S)$$

Henceforth, we consider $(\pi_{\lambda}, V_{\lambda})$ over $\bar{K}_1$ until the end of this section. Assume that $\lambda' = \lambda$. We define $\text{STab}(\lambda)^b$ and $\text{STab}(\lambda)^{-b}$ to be the sets of all the standard tableaux T of shape λ such that the smallest number which is not assigned in diagonals of Young diagrams appears in the first row and the first column respectively. Both $\text{STab}(\lambda)^b$ and $\text{STab}(\lambda)^{-b}$ are not empty if $\lambda \neq (1, 1)$. For $\lambda \neq (1, 1)$, $\text{STab}(\lambda)$ is the disjoint union of $\text{STab}(\lambda)^b$ and $\text{STab}(\lambda)^{-b}$. There exists an involutive bijection from $\text{STab}(\lambda)^b$ to $\text{STab}(\lambda)^{-b}$ which is defined by $T \mapsto T'$. Now, we define two subspaces $V_{\lambda, b}^+$ and $V_{\lambda, b}^-$ of V_{λ} as follows.

$$\begin{aligned} V_{\lambda, b}^+ &= \bigoplus_{T \in \text{STab}(\lambda)^b} \bar{K}_1(\sqrt{\psi(T')}v_T + \sqrt{\psi(T)}v_{T'}) \\ V_{\lambda, b}^- &= \bigoplus_{T \in \text{STab}(\lambda)^b} \bar{K}_1(\sqrt{\psi(T')}v_T - \sqrt{\psi(T)}v_{T'}) \end{aligned}$$

Then, it is clear that $\dim_{\bar{K}_1} V_{\lambda, b}^+ = \dim_{\bar{K}_1} V_{\lambda, b}^- = \dim_{\bar{K}_1} V_{\lambda, b}/2$ and $V_{\lambda, b}$ is a direct sum of $V_{\lambda, b}^+$ and $V_{\lambda, b}^-$ as vector spaces over $\bar{K}_1$.

Proposition 6.12. *If $\lambda' = \lambda$ and $\lambda \neq (1, 1)$, then $V_{\lambda, b} = V_{\lambda, b}^+ \oplus V_{\lambda, b}^-$ is a direct sum decomposition as $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^b$ -submodules.*

Proof. For $T \in \text{STab}(\lambda)^b$, we have from Corollary 6.11

$$\begin{aligned} & \pi_{\lambda, b}(b) \left(\sqrt{\psi(T')}v_T \pm \sqrt{\psi(T)}v_{T'} \right) \\ &= \sum_{S \in \text{STab}(\lambda)} \left\{ \left(\pi_{\lambda, b}(b) \right)_{T, S} \sqrt{\psi(T')}v_S \pm \left(\pi_{\lambda, b}(b) \right)_{T', S'} \sqrt{\psi(T)}v_{S'} \right\} \\ &= \sum_{S \in \text{STab}(\lambda)} \sqrt{\left(\pi_{\lambda, b}(b) \right)_{T, S} \left(\pi_{\lambda, b}(b) \right)_{T', S'}} \left(\sqrt{\psi(S')}v_S \pm \sqrt{\psi(S)}v_{S'} \right) \\ &= \sum_{S \in \text{STab}(\lambda)^b} \left\{ \sqrt{\left(\pi_{\lambda, b}(b) \right)_{T, S} \left(\pi_{\lambda, b}(b) \right)_{T', S'}} \pm \sqrt{\left(\pi_{\lambda, b}(b) \right)_{T, S'} \left(\pi_{\lambda, b}(b) \right)_{T', S}} \right\} \\ & \quad \times \left(\sqrt{\psi(S')}v_S \pm \sqrt{\psi(S)}v_{S'} \right) \end{aligned}$$

Thus, $\pi_{\lambda, b}(b) \left(\sqrt{\psi(T')}v_T \pm \sqrt{\psi(T)}v_{T'} \right)$ is in $V_{\lambda, b}^{\pm}$. □

If $\lambda = (1, 1)$, then $T' = T$. Moreover $s_1 T = T^*$ for $T \in \text{STab}(\lambda)$. Since axial distances $d_{T, 2, 1} = 0$ for all $T \in \text{STab}(\lambda)$ in this case, (b2)-(c) and (e2)-(c) reduce to

$$\pi_{\lambda}(b_2) \langle v_T, v_{s_1 T} \rangle = \langle v_T, v_{s_1 T} \rangle \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad \pi_{\lambda'}(b_2) \langle v_{T'}, v_{s_1 T'} \rangle = \langle v_{T'}, v_{s_1 T'} \rangle \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

So letting $T_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, we have two submodules $V_{\lambda,b}^+ = \bar{K}_1 v_{T_1}$, $V_{\lambda,b}^- = \bar{K}_1 v_{T_1^*}$. One can readily see that $V_{\lambda,b} = V_{\lambda,b}^+ \oplus V_{\lambda,b}^-$ as $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^b$ -submodules.

Theorem 6.13. (1) $\pi_{\lambda',b} \cong \pi_{\lambda,b}$.

(2) If $\lambda' = \lambda$, then $\pi_{\lambda,b}$ decomposes into inequivalent subrepresentations $\pi_{\lambda,b}^+$ and $\pi_{\lambda,b}^-$ over $\bar{K}_1$, the algebraic closure of K_1 . Furthermore, $\deg \pi_{\lambda,b}^+ = \deg \pi_{\lambda,b}^- = \deg \pi_{\lambda,b}/2$.

(3) Let $\{\lambda_i, \lambda'_i, \mu_j\}_{i,j}$ be the set of 2-tuples of Young diagrams such that

$$\lambda'_i \neq \lambda_i, \quad \mu'_j = \mu_j$$

Then

$$\text{Irr}(\mathcal{H}_{\bar{K}_1, B_n}(q)^b) = \{\pi_{\lambda_i,b}, \pi_{\mu_j,b}^+, \pi_{\mu_j,b}^-\}_{i,j}$$

is a basic set of irreducible representations of $\mathcal{H}_{\bar{K}_1, B_n}(q)^b$. $\mathcal{H}_{\bar{K}_1, B_n}(q)^b$ is semisimple.

Proof. (1) is Corollary 6.10 itself. The submodules $V_{\lambda,b}^+$ and $V_{\lambda,b}^-$ correspond to the case (2) of Theorem 2.7. Hence they are simple and mutually non-isomorphic. Thus (2) holds. The proof of (3) is in the same manner as in the proof of Theorem 6.2, just replacing $\natural$ and $\bar{K}_0$ with b and $\bar{K}_1$ respectively. So we omit the detail. $\square$

7 The branching rule for the Hecke algebras of Type D

Henceforth we assume $n \geq 4$. We denote by $(\pi_{\lambda,\dagger}, V_{\lambda,\dagger})$, $(\pi_{\lambda,\natural,\dagger}, V_{\lambda,\natural,\dagger})$, $(\pi_{\lambda,\sharp,\dagger}, V_{\lambda,\sharp,\dagger})$, $(\pi_{\lambda,b,\dagger}, V_{\lambda,b,\dagger})$ the restrictions of (π_λ, V_λ) , $(\pi_{\lambda,\natural}, V_{\lambda,\natural})$, $(\pi_{\lambda,\sharp}, V_{\lambda,\sharp})$, $(\pi_{\lambda,b}, V_{\lambda,b})$ to $\mathcal{H}_{K_1, B_n}(1, q)^\dagger$ respectively. It is clear that $\pi_{\lambda,\dagger} \cong \pi_{\lambda,\natural,\dagger} \cong \pi_{\lambda,\sharp,\dagger} \cong \pi_{\lambda,b,\dagger}$. If $\lambda^* = \lambda$, then we denote by $(\pi_{\lambda,\sharp,\dagger}^\pm, V_{\lambda,\sharp,\dagger}^\pm)$ the restriction of $(\pi_{\lambda,\sharp}^\pm, V_{\lambda,\sharp}^\pm)$ to $\mathcal{H}_{K_1, B_n}(1, q)^\dagger$. We adopt the similar notations for the cases $\lambda' = \lambda$ and $\lambda'^* = \lambda$. We notice that $\mathcal{H}_{K_1, B_n}(1, q)^\sharp = \mathcal{H}_{K_1, D_n}(q)$ and $\mathcal{H}_{K_1, B_n}(1, q)^b$ and $\mathcal{H}_{K_1, B_n}(1, q)^\natural$ are all $\mathbb{Z}_2$ -crossed products with $A_{\bar{0}} = \mathcal{H}_{K_1, B_n}(1, q)^\dagger$.

Proposition 7.1. $\pi_{\lambda,\dagger} \cong \pi_{\lambda^*,\dagger} \cong \pi_{\lambda',\dagger} \cong \pi_{\lambda'^*,\dagger}$

Proof. Since $\mathcal{H}_{K_1, B_n}(1, q)^\dagger = \mathcal{H}_{K_1, B_n}(1, q)^\sharp \cap \mathcal{H}_{K_1, B_n}(1, q)^b \cap \mathcal{H}_{K_1, B_n}(1, q)^\natural$, we have that $\pi_{\lambda,\dagger}, \pi_{\lambda^*,\dagger}, \pi_{\lambda',\dagger}$ and $\pi_{\lambda'^*,\dagger}$ are all equivalent. $\square$

All the representations $V_\lambda, V_{\lambda,\sharp}, V_{\lambda,b}, V_{\lambda,\natural}, \dots$ can be defined over $\bar{K}_1$, so we consider all these over $\bar{K}_1$. At first, we assume that $\lambda, \lambda^*, \lambda'$ and λ'^* are mutually different.

Proposition 7.2. Let $\lambda, \lambda^*, \lambda'$ and λ'^* are mutually different. Then $\pi_{\lambda,\sharp,\dagger}$ is irreducible.

Proof. We have already shown that $\pi_{\lambda,\sharp}$ and $\pi_{\lambda',\sharp}$ are irreducible and inequivalent. On the other hand, $\pi_{\lambda,\sharp,\dagger} \cong \pi_{\lambda',\sharp,\dagger}$ by Proposition 7.1. This corresponds to the case (1) of Theorem 2.7. Thus we can conclude that $\pi_{\lambda,\sharp,\dagger}$ is irreducible. $\square$

Next, we assume that only one of the three cases $\lambda^* = \lambda$, $\lambda' = \lambda$, $\lambda'^* = \lambda$ holds, and considering representations over $\bar{K}_1$, the algebraic closure of K_1

Proposition 7.3.

- (1) If $\lambda^* = \lambda$, $\lambda' \neq \lambda$, $\lambda'^* \neq \lambda$, then $\pi_{\lambda, \#}^+ \cong \pi_{\lambda', \#}^+$ and $\pi_{\lambda, \#}^- \cong \pi_{\lambda', \#}^-$. They are irreducible and mutually inequivalent.
- (2) If $\lambda^* \neq \lambda$, $\lambda' = \lambda$, $\lambda'^* \neq \lambda$, then $\pi_{\lambda, \#}^+ \cong \pi_{\lambda', \#}^+$ and $\pi_{\lambda, \#}^- \cong \pi_{\lambda', \#}^-$. They are irreducible and mutually inequivalent.
- (3) If $\lambda^* \neq \lambda$, $\lambda' \neq \lambda$, $\lambda'^* = \lambda$, then $\pi_{\lambda, \#}^+ \cong \pi_{\lambda', \#}^+$ and $\pi_{\lambda, \#}^- \cong \pi_{\lambda', \#}^-$. They are irreducible and mutually inequivalent.

Proof. When $\lambda^* = \lambda$, $\lambda'^* = \lambda'$ also holds. We have already shown that $\pi_{\lambda, \#}$ and $\pi_{\lambda', \#}$ have the irreducible decompositions $\pi_{\lambda, \#} = \pi_{\lambda, \#}^+ \oplus \pi_{\lambda, \#}^-$ and $\pi_{\lambda', \#} = \pi_{\lambda', \#}^+ \oplus \pi_{\lambda', \#}^-$ respectively. $\pi_{\lambda, \#} \not\cong \pi_{\lambda', \#}$ since $\lambda' \neq \lambda$, while $\pi_{\lambda, \#} \cong \pi_{\lambda', \#}$. Therefore, $V_{\lambda, \#}^+$ is $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger$ -isomorphic to a submodule of $V_{\lambda', \#}^+$. This corresponds to the case (1) of Theorem 2.7, hence $\pi_{\lambda, \#}^+$ is irreducible. In the same manner, $\pi_{\lambda, \#}^+$, $\pi_{\lambda', \#}^+$, $\pi_{\lambda, \#}^-$, $\pi_{\lambda', \#}^-$ are also irreducible. Assume $\pi_{\lambda, \#}^+ \cong \pi_{\lambda', \#}^-$. Then four irreducible representations $\pi_{\lambda, \#}^\pm$, $\pi_{\lambda', \#}^\pm$ are all equivalent. But this is impossible because of Theorem 2.7. Thus $\pi_{\lambda, \#}^+ \not\cong \pi_{\lambda', \#}^-$ holds. Consequently, we also have $\pi_{\lambda, \#}^+ \cong \pi_{\lambda', \#}^+$ and $\pi_{\lambda, \#}^- \cong \pi_{\lambda', \#}^-$. In the same manner, we also obtain (2) and (3). $\square$

Last, we assume $\lambda^* = \lambda$ and $\lambda' = \lambda$ (these imply $\lambda'^* = \lambda$). For brevity, we set

$$\begin{aligned} v_T^{++} &= \sqrt[4]{\psi(T')\psi(T'^*)}(v_T + v_{T^*}) + \sqrt[4]{\psi(T)\psi(T^*)}(v_{T'} + v_{T'^*}), \\ v_T^{+-} &= \sqrt[4]{\psi(T')\psi(T'^*)}(v_T + v_{T^*}) - \sqrt[4]{\psi(T)\psi(T^*)}(v_{T'} + v_{T'^*}), \\ v_T^{-+} &= \sqrt[4]{\psi(T')\psi(T'^*)}(v_T - v_{T^*}) + \sqrt[4]{\psi(T)\psi(T^*)}(v_{T'} - v_{T'^*}), \\ v_T^{--} &= \sqrt[4]{\psi(T')\psi(T'^*)}(v_T - v_{T^*}) - \sqrt[4]{\psi(T)\psi(T^*)}(v_{T'} - v_{T'^*}). \end{aligned}$$

We define $\bar{K}_1$ -subspaces $V_{\lambda, \#}^{++}$, $V_{\lambda, \#}^{+-}$, $V_{\lambda, \#}^{-+}$, $V_{\lambda, \#}^{--}$ of V_λ as follows.

$$\begin{aligned} V_{\lambda, \#}^{++} &= \bigoplus_{T \in \text{STab}(\lambda)^\dagger} \bar{K}_1 v_T^{++}, & V_{\lambda, \#}^{+-} &= \bigoplus_{T \in \text{STab}(\lambda)^\dagger} \bar{K}_1 v_T^{+-}, \\ V_{\lambda, \#}^{-+} &= \bigoplus_{T \in \text{STab}(\lambda)^\dagger} \bar{K}_1 v_T^{-+}, & V_{\lambda, \#}^{--} &= \bigoplus_{T \in \text{STab}(\lambda)^\dagger} \bar{K}_1 v_T^{--} \end{aligned}$$

where $\text{STab}(\lambda)^\dagger = \text{STab}(\lambda)^\# \cap \text{STab}(\lambda)^\flat$.

Remark 7.4. We notice that each fourth square root has four branches. We take a suitable branch in each computation of fourth square roots for consistency.

Proposition 7.5.

$$V_{\lambda, \#}^{++} \subseteq V_{\lambda, \#}^+ \cap V_{\lambda, \#}^+, \quad V_{\lambda, \#}^{+-} \subseteq V_{\lambda, \#}^+ \cap V_{\lambda, \#}^-, \quad V_{\lambda, \#}^{-+} \subseteq V_{\lambda, \#}^- \cap V_{\lambda, \#}^+, \quad V_{\lambda, \#}^{--} \subseteq V_{\lambda, \#}^- \cap V_{\lambda, \#}^-,$$

and furthermore, we have direct sum decompositions as $\bar{K}_1$ -vector spaces as follows.

$$\begin{aligned} V_{\lambda, \sharp}^+ &= V_{\lambda, \sharp, \dagger}^{++} \oplus V_{\lambda, \sharp, \dagger}^{+-}, & V_{\lambda, \sharp}^- &= V_{\lambda, \sharp, \dagger}^{-+} \oplus V_{\lambda, \sharp, \dagger}^{--}, \\ V_{\lambda, b}^+ &= V_{\lambda, \sharp, \dagger}^{++} \oplus V_{\lambda, \sharp, \dagger}^{+-}, & V_{\lambda, b}^- &= V_{\lambda, \sharp, \dagger}^{-+} \oplus V_{\lambda, \sharp, \dagger}^{--}, \\ V_{\lambda, \natural}^+ &= V_{\lambda, \sharp, \dagger}^{++} \oplus V_{\lambda, \sharp, \dagger}^{+-}, & V_{\lambda, \natural}^- &= V_{\lambda, \sharp, \dagger}^{-+} \oplus V_{\lambda, \sharp, \dagger}^{--}. \end{aligned}$$

Proof. Assume $T \in \text{STab}(\lambda)^\dagger$. Then T and T' belong to $\text{STab}(\lambda)^\sharp$. Therefore, we have $V_{\lambda, \sharp, \dagger}^{++}, V_{\lambda, \sharp, \dagger}^{+-} \in V_{\lambda, \sharp}^+$ and $V_{\lambda, \sharp, \dagger}^{-+}, V_{\lambda, \sharp, \dagger}^{--} \in V_{\lambda, \sharp}^-$ immediately. We also see that $T, T^* \in \text{STab}(\lambda)^\flat$ and $T, T' \in \text{STab}(\lambda)^\natural$. If $n/2$ is even, then $\psi(T^*) = \psi(T)$ and hence we have two expressions for each linear combination as follows.

$$\begin{aligned} v_T^{++} &= \begin{cases} \sqrt{\psi(T')}v_T + \sqrt{\psi(T)}v_{T'} + \sqrt{\psi(T'^*)}v_{T^*} + \sqrt{\psi(T^*)}v_{T'^*} \in V_{\lambda, b}^+, \\ \sqrt{\psi(T'^*)}v_T + \sqrt{\psi(T)}v_{T'^*} + \sqrt{\psi(T^*)}v_{T'} + \sqrt{\psi(T')}v_{T^*} \in V_{\lambda, \natural}^+, \end{cases} \\ v_T^{+-} &= \begin{cases} \sqrt{\psi(T')}v_T - \sqrt{\psi(T)}v_{T'} + \sqrt{\psi(T'^*)}v_{T^*} - \sqrt{\psi(T^*)}v_{T'^*} \in V_{\lambda, b}^-, \\ \sqrt{\psi(T'^*)}v_T - \sqrt{\psi(T)}v_{T'^*} - (\sqrt{\psi(T^*)}v_{T'} - \sqrt{\psi(T')}v_{T^*}) \in V_{\lambda, \natural}^-, \end{cases} \\ v_T^{-+} &= \begin{cases} \sqrt{\psi(T')}v_T + \sqrt{\psi(T)}v_{T'} - (\sqrt{\psi(T'^*)}v_{T^*} + \sqrt{\psi(T^*)}v_{T'^*}) \in V_{\lambda, b}^+, \\ \sqrt{\psi(T'^*)}v_T - \sqrt{\psi(T)}v_{T'^*} + \sqrt{\psi(T^*)}v_{T'} - \sqrt{\psi(T')}v_{T^*} \in V_{\lambda, \natural}^-, \end{cases} \\ v_T^{--} &= \begin{cases} \sqrt{\psi(T')}v_T - \sqrt{\psi(T)}v_{T'} - (\sqrt{\psi(T'^*)}v_{T^*} - \sqrt{\psi(T^*)}v_{T'^*}) \in V_{\lambda, b}^-, \\ \sqrt{\psi(T'^*)}v_T + \sqrt{\psi(T)}v_{T'^*} - (\sqrt{\psi(T^*)}v_{T'} + \sqrt{\psi(T')}v_{T^*}) \in V_{\lambda, \natural}^+. \end{cases} \end{aligned}$$

If $n/2$ is odd, then $\psi(T^*) = -\psi(T)$ and hence

$$\begin{aligned} v_T^{++} &= \begin{cases} \sqrt[4]{-1}\{\sqrt{\psi(T')}v_T + \sqrt{\psi(T)}v_{T'} + \sqrt{\psi(T'^*)}v_{T^*} + \sqrt{\psi(T^*)}v_{T'^*}\} \in V_{\lambda, b}^+, \\ \sqrt[4]{-1}\{\sqrt{\psi(T'^*)}v_T + \sqrt{\psi(T)}v_{T'^*} + \sqrt{\psi(T^*)}v_{T'} + \sqrt{\psi(T')}v_{T^*}\} \in V_{\lambda, \natural}^+, \end{cases} \\ v_T^{+-} &= \begin{cases} \sqrt[4]{-1}\{\sqrt{\psi(T')}v_T - \sqrt{\psi(T)}v_{T'} + \sqrt{\psi(T'^*)}v_{T^*} - \sqrt{\psi(T^*)}v_{T'^*}\} \in V_{\lambda, b}^-, \\ \sqrt[4]{-1}\{\sqrt{\psi(T'^*)}v_T - \sqrt{\psi(T)}v_{T'^*} - (\sqrt{\psi(T^*)}v_{T'} - \sqrt{\psi(T')}v_{T^*})\} \in V_{\lambda, \natural}^-, \end{cases} \\ v_T^{-+} &= \begin{cases} \sqrt[4]{-1}\{\sqrt{\psi(T')}v_T + \sqrt{\psi(T)}v_{T'} - (\sqrt{\psi(T'^*)}v_{T^*} + \sqrt{\psi(T^*)}v_{T'^*})\} \in V_{\lambda, b}^+, \\ \sqrt[4]{-1}\{\sqrt{\psi(T'^*)}v_T - \sqrt{\psi(T)}v_{T'^*} + \sqrt{\psi(T^*)}v_{T'} - \sqrt{\psi(T')}v_{T^*}\} \in V_{\lambda, \natural}^-, \end{cases} \\ v_T^{--} &= \begin{cases} \sqrt[4]{-1}\{\sqrt{\psi(T')}v_T - \sqrt{\psi(T)}v_{T'} - (\sqrt{\psi(T'^*)}v_{T^*} - \sqrt{\psi(T^*)}v_{T'^*})\} \in V_{\lambda, b}^-, \\ \sqrt[4]{-1}\{\sqrt{\psi(T'^*)}v_T + \sqrt{\psi(T)}v_{T'^*} - (\sqrt{\psi(T^*)}v_{T'} + \sqrt{\psi(T')}v_{T^*})\} \in V_{\lambda, \natural}^+. \end{cases} \end{aligned}$$

Thus the first assertion holds. We also obtain that

$$\begin{aligned} v_T^{++} + v_T^{+-} &= 2\sqrt[4]{\psi(T')\psi(T'^*)}(v_T + v_{T^*}), \\ v_T^{++} - v_T^{+-} &= 2\sqrt[4]{\psi(T)\psi(T^*)}(v_{T'} + v_{T'^*}). \end{aligned}$$

Thus every $v_T + v_{T^*}$ ($T \in \text{STab}(\lambda)^\sharp$) belongs to $V_{\lambda, \sharp, \dagger}^{++} + V_{\lambda, \sharp, \dagger}^{+-}$. We readily see that $V_{\lambda, \sharp, \dagger}^{++} + V_{\lambda, \sharp, \dagger}^{+-}$ is a direct sum by the first assertion. Therefore we obtain $V_{\lambda, \sharp}^+ = V_{\lambda, \sharp, \dagger}^{++} \oplus V_{\lambda, \sharp, \dagger}^{+-}$. The others are obtained in the same manner. $\square$

Proposition 7.6. $V_{\lambda, \sharp, \dagger}^{++}, V_{\lambda, \sharp, \dagger}^{+-}, V_{\lambda, \sharp, \dagger}^{-+}, V_{\lambda, \sharp, \dagger}^{--}$ are $\mathcal{H}_{\tilde{K}_1, B_n}(1, q)^\dagger$ -modules.

Proof. Using Corollary 6.5 and Corollary 6.11, one can check by a direct computation that

$$\begin{aligned}
& \pi_\lambda(b) \left(\sqrt[4]{\psi(T')\psi(T'^*)} v_{T'} \right) \\
&= \sum_{S \in \text{STab}(\lambda)^\dagger} \left\{ \sqrt[4]{(\pi_\lambda(b))_{TS} (\pi_\lambda(b))_{T^*S^*} (\pi_\lambda(b))_{T'S'} (\pi_\lambda(b))_{T'^*S'^*}} \sqrt[4]{\psi(S')\psi(S'^*)} v_S \right. \\
&+ \sqrt[4]{(\pi_\lambda(b))_{TS'} (\pi_\lambda(b))_{T^*S'^*} (\pi_\lambda(b))_{T'S} (\pi_\lambda(b))_{T'^*S^*}} \sqrt[4]{\psi(S)\psi(S^*)} v_{S'} \\
&+ \sqrt[4]{(\pi_\lambda(b))_{TS^*} (\pi_\lambda(b))_{T^*S} (\pi_\lambda(b))_{T'S'^*} (\pi_\lambda(b))_{T'^*S'}} \sqrt[4]{\psi(S'^*)\psi(S')} v_{S^*} \\
&\left. + \sqrt[4]{(\pi_\lambda(b))_{TS'^*} (\pi_\lambda(b))_{T^*S'} (\pi_\lambda(b))_{T'S^*} (\pi_\lambda(b))_{T'^*S}} \sqrt[4]{\psi(S^*)\psi(S)} v_{S'^*} \right\}
\end{aligned}$$

for $b \in \mathcal{H}_{\tilde{K}_1, B_n}(1, q)^\dagger$. Similarly we have

$$\begin{aligned}
& \pi_\lambda(b) \left(\sqrt[4]{\psi(T)\psi(T^*)} v_{T'} \right) \\
&= \sum_{S \in \text{STab}(\lambda)^\dagger} \left\{ \sqrt[4]{(\pi_\lambda(b))_{T'S} (\pi_\lambda(b))_{T'^*S^*} (\pi_\lambda(b))_{TS'} (\pi_\lambda(b))_{T^*S'^*}} \sqrt[4]{\psi(S')\psi(S'^*)} v_S \right. \\
&+ \sqrt[4]{(\pi_\lambda(b))_{T'S'} (\pi_\lambda(b))_{T'^*S'^*} (\pi_\lambda(b))_{TS} (\pi_\lambda(b))_{T^*S^*}} \sqrt[4]{\psi(S)\psi(S^*)} v_{S'} \\
&+ \sqrt[4]{(\pi_\lambda(b))_{T'S^*} (\pi_\lambda(b))_{T'^*S} (\pi_\lambda(b))_{TS'^*} (\pi_\lambda(b))_{T^*S'}} \sqrt[4]{\psi(S'^*)\psi(S')} v_{S^*} \\
&\left. + \sqrt[4]{(\pi_\lambda(b))_{T'S'^*} (\pi_\lambda(b))_{T'^*S'} (\pi_\lambda(b))_{TS^*} (\pi_\lambda(b))_{T^*S}} \sqrt[4]{\psi(S^*)\psi(S)} v_{S'^*} \right\} \\
& \pi_\lambda(b) \left(\sqrt[4]{\psi(T'^*)\psi(T')} v_{T^*} \right) \\
&= \sum_{S \in \text{STab}(\lambda)^\dagger} \left\{ \sqrt[4]{(\pi_\lambda(b))_{T^*S} (\pi_\lambda(b))_{TS^*} (\pi_\lambda(b))_{T'^*S'} (\pi_\lambda(b))_{T'^*S'^*}} \sqrt[4]{\psi(S')\psi(S'^*)} v_S \right. \\
&+ \sqrt[4]{(\pi_\lambda(b))_{T^*S'} (\pi_\lambda(b))_{TS'^*} (\pi_\lambda(b))_{T'^*S} (\pi_\lambda(b))_{T'^*S^*}} \sqrt[4]{\psi(S)\psi(S^*)} v_{S'} \\
&+ \sqrt[4]{(\pi_\lambda(b))_{T^*S^*} (\pi_\lambda(b))_{TS} (\pi_\lambda(b))_{T'^*S'^*} (\pi_\lambda(b))_{T'^*S'}} \sqrt[4]{\psi(S'^*)\psi(S')} v_{S^*} \\
&\left. + \sqrt[4]{(\pi_\lambda(b))_{T^*S'^*} (\pi_\lambda(b))_{TS'} (\pi_\lambda(b))_{T'^*S^*} (\pi_\lambda(b))_{T'^*S}} \sqrt[4]{\psi(S^*)\psi(S)} v_{S'^*} \right\} \\
& \pi_\lambda(b) \left(\sqrt[4]{\psi(T')\psi(T'^*)} v_{T'^*} \right) \\
&= \sum_{S \in \text{STab}(\lambda)^\dagger} \left\{ \sqrt[4]{(\pi_\lambda(b))_{T'^*S} (\pi_\lambda(b))_{T'^*S^*} (\pi_\lambda(b))_{T^*S'} (\pi_\lambda(b))_{T^*S'^*}} \sqrt[4]{\psi(S')\psi(S'^*)} v_S \right. \\
&+ \sqrt[4]{(\pi_\lambda(b))_{T'^*S'} (\pi_\lambda(b))_{T'^*S'^*} (\pi_\lambda(b))_{T^*S} (\pi_\lambda(b))_{T^*S^*}} \sqrt[4]{\psi(S)\psi(S^*)} v_{S'} \\
&+ \sqrt[4]{(\pi_\lambda(b))_{T'^*S^*} (\pi_\lambda(b))_{T'^*S} (\pi_\lambda(b))_{T^*S'^*} (\pi_\lambda(b))_{T^*S'}} \sqrt[4]{\psi(S'^*)\psi(S')} v_{S^*} \\
&\left. + \sqrt[4]{(\pi_\lambda(b))_{T'^*S'^*} (\pi_\lambda(b))_{T'^*S'} (\pi_\lambda(b))_{T^*S^*} (\pi_\lambda(b))_{T^*S}} \sqrt[4]{\psi(S^*)\psi(S)} v_{S'^*} \right\}
\end{aligned}$$

Summing up all of those, we get

$$\begin{aligned} \pi_\lambda(b)v_T^{++} = & \sum_{S \in \text{STab}(\lambda)^\dagger} \left\{ \sqrt[4]{(\pi_\lambda(b))_{TS}(\pi_\lambda(b))_{T^*S^*}(\pi_\lambda(b))_{T'S'}(\pi_\lambda(b))_{T'^*S'^*}} \right. \\ & + \sqrt[4]{(\pi_\lambda(b))_{T'S}(\pi_\lambda(b))_{T'^*S^*}(\pi_\lambda(b))_{TS'}(\pi_\lambda(b))_{T^*S'^*}} \\ & + \sqrt[4]{(\pi_\lambda(b))_{T^*S}(\pi_\lambda(b))_{TS^*}(\pi_\lambda(b))_{T'^*S'}(\pi_\lambda(b))_{T'S'^*}} \\ & \left. + \sqrt[4]{(\pi_\lambda(b))_{T'^*S}(\pi_\lambda(b))_{T'S^*}(\pi_\lambda(b))_{T^*S'}(\pi_\lambda(b))_{TS'^*}} \right\} v_S^{++}. \end{aligned}$$

Thus $\pi_\lambda(b)V_{\lambda, \sharp, \dagger}^{++} \subseteq V_{\lambda, \sharp, \dagger}^{++}$ for $b \in \mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger$. In the same fashion, we also obtain $\pi_\lambda(b)V_{\lambda, \sharp, \dagger}^{+-} \subseteq V_{\lambda, \sharp, \dagger}^{+-}$, $\pi_\lambda(b)V_{\lambda, \sharp, \dagger}^{-+} \subseteq V_{\lambda, \sharp, \dagger}^{-+}$ and $\pi_\lambda(b)V_{\lambda, \sharp, \dagger}^{--} \subseteq V_{\lambda, \sharp, \dagger}^{--}$. $\square$

We denote the representations of $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger$ given in Proposition 7.6 by $(\pi_{\lambda, \sharp, \dagger}^{++}, V_{\lambda, \sharp, \dagger}^{++})$, $(\pi_{\lambda, \sharp, \dagger}^{+-}, V_{\lambda, \sharp, \dagger}^{+-})$, $(\pi_{\lambda, \sharp, \dagger}^{-+}, V_{\lambda, \sharp, \dagger}^{-+})$ and $(\pi_{\lambda, \sharp, \dagger}^{--}, V_{\lambda, \sharp, \dagger}^{--})$.

Proposition 7.7. $(\pi_{\lambda, \sharp, \dagger}^{++}, V_{\lambda, \sharp, \dagger}^{++})$, $(\pi_{\lambda, \sharp, \dagger}^{+-}, V_{\lambda, \sharp, \dagger}^{+-})$, $(\pi_{\lambda, \sharp, \dagger}^{-+}, V_{\lambda, \sharp, \dagger}^{-+})$, $(\pi_{\lambda, \sharp, \dagger}^{--}, V_{\lambda, \sharp, \dagger}^{--})$ are mutually non-isomorphic simple left $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger$ -modules.

Proof. Observing that $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\sharp$ and $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\flat$ and $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\natural$ are all $\mathbb{Z}_2$ -crossed products with $A_{\bar{0}} = \mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger$, one can deduce the assertion directly from Proposition 7.5, Proposition 7.6 and Theorem 2.7 (2). $\square$

We have the following immediately from Proposition 7.5, Proposition 7.6 and Proposition 7.7.

Corollary 7.8. We have branching rules for λ such that $\lambda = \lambda^* = \lambda' = \lambda'^*$ as follows.

$$\begin{aligned} \pi_{\lambda, \dagger} &= \pi_{\lambda, \sharp, \dagger}^{++} \oplus \pi_{\lambda, \sharp, \dagger}^{+-} \oplus \pi_{\lambda, \sharp, \dagger}^{-+} \oplus \pi_{\lambda, \sharp, \dagger}^{--}, \\ \pi_{\lambda, \sharp, \dagger}^+ &= \pi_{\lambda, \sharp, \dagger}^{++} \oplus \pi_{\lambda, \sharp, \dagger}^{+-}, & \pi_{\lambda, \sharp, \dagger}^- &= \pi_{\lambda, \sharp, \dagger}^{-+} \oplus \pi_{\lambda, \sharp, \dagger}^{--}, \\ \pi_{\lambda, \flat, \dagger}^+ &= \pi_{\lambda, \sharp, \dagger}^{++} \oplus \pi_{\lambda, \sharp, \dagger}^{-+}, & \pi_{\lambda, \flat, \dagger}^- &= \pi_{\lambda, \sharp, \dagger}^{+-} \oplus \pi_{\lambda, \sharp, \dagger}^{--}, \\ \pi_{\lambda, \natural, \dagger}^+ &= \pi_{\lambda, \sharp, \dagger}^{++} \oplus \pi_{\lambda, \sharp, \dagger}^{--}, & \pi_{\lambda, \natural, \dagger}^- &= \pi_{\lambda, \sharp, \dagger}^{+-} \oplus \pi_{\lambda, \sharp, \dagger}^{-+}, \end{aligned}$$

$\pi_{\lambda, \sharp, \dagger}^{++}$, $\pi_{\lambda, \sharp, \dagger}^{+-}$, $\pi_{\lambda, \sharp, \dagger}^{-+}$, $\pi_{\lambda, \sharp, \dagger}^{--}$ are irreducible and mutually inequivalent. They have the same degree.

Theorem 7.9. Let $\{\lambda_i, \lambda_i^*, \lambda_i', \lambda_i'^*, \mu_j, \mu_j', \nu_k, \nu_k^*, \nu_k', \nu_k'^*, \kappa_l, \kappa_l', \kappa_l^*, \kappa_l'^*, \iota_m, \iota_m^*, \iota_m', \iota_m'^*\}_{i,j,k,l,m}$ be the set of 2-tuples of Young diagrams such that

$$\begin{aligned} \lambda_i' &\neq \lambda_i, \lambda_i^* \neq \lambda_i, \lambda_i'^* \neq \lambda_i, & \mu_j^* &= \mu_j, \mu_j' \neq \mu_j, \mu_j'^* \neq \mu_j, \\ \nu_k^* &\neq \nu_k, \nu_k' = \nu_k, \nu_k'^* \neq \nu_k, & \kappa_l^* &\neq \kappa_l, \kappa_l' \neq \kappa_l, \kappa_l'^* = \kappa_l, & \iota_m' &= \iota_m^* = \iota_m'^* = \iota_m. \end{aligned}$$

Then

$$\text{Irr}(\mathcal{H}_{\bar{K}_1, B_n}(q)^\dagger) = \{\pi_{\lambda_i, \dagger}, \pi_{\mu_j, \sharp, \dagger}^\pm, \pi_{\nu_k, \flat, \dagger}^\pm, \pi_{\kappa_l, \natural, \dagger}^\pm, \pi_{\iota_m, \dagger}^{++}, \pi_{\iota_m, \dagger}^{+-}, \pi_{\iota_m, \dagger}^{-+}, \pi_{\iota_m, \dagger}^{--}\}_{i,j,k,l,m}$$

is a basic set of irreducible representations of $\mathcal{H}_{\bar{K}_1, B_n}(q)^\dagger$. Moreover $\mathcal{H}_{\bar{K}_1, B_n}(q)^\dagger$ is semisimple.

Proof. We show that the elements of $\text{Irr}(\mathcal{H}_{\bar{K}_1, B_n}(q)^\dagger)$ are mutually inequivalent. We have already proved (1)–(4) below.

- (1) $\pi_{\mu_j, \#}^+ \not\cong \pi_{\mu_j, \#}^-$, (2) $\pi_{\nu_k, \#}^+ \not\cong \pi_{\nu_k, \#}^-$, (3) $\pi_{\kappa_l, \#}^+ \not\cong \pi_{\kappa_l, \#}^-$,
(4) $\pi_{\iota_m, \#}^{++}, \pi_{\iota_m, \#}^{+-}, \pi_{\iota_m, \#}^{-+}, \pi_{\iota_m, \#}^{--}$ are mutually inequivalent.

Indeed, (1)–(3) have proved in Proposition 7.3, and (4) in Proposition 7.7. For brevity, we denote $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\#$ by $\mathcal{H}^\#$ and $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger$ by $\mathcal{H}^\dagger$, and so on. We shall give inductions of elements of $\text{Irr}(\mathcal{H}_{\bar{K}_1, B_n}(q)^\dagger)$ to $\mathcal{H} = \mathcal{H}_{\bar{K}_1, B_n}(1, q)$. We consider five cases depending upon the form of the 2-tuple of Young diagrams.

case 1 : $\lambda_i' \neq \lambda_i, \lambda_i^* \neq \lambda_i, \lambda_i'^* \neq \lambda_i$

In this case, the induction of $V_{\lambda_i, \#}^\dagger$ to $\mathcal{H}^\#$ yields $(V_{\lambda_i, \#}^\dagger)^{\mathcal{H}^\#} \cong V_{\lambda_i, \#} \oplus V_{\lambda_i', \#}$ by Theorem 2.7 (1). Moreover, the induction of $V_{\lambda_i, \#}$ to $\mathcal{H}$ yields $(V_{\lambda_i, \#})^{\mathcal{H}} \cong V_{\lambda_i} \oplus V_{\lambda_i^*}$ by Theorem 2.7 (1) again. Thus we obtain $(V_{\lambda_i, \#}^\dagger)^{\mathcal{H}} \cong V_{\lambda_i} \oplus V_{\lambda_i^*} \oplus V_{\lambda_i'} \oplus V_{\lambda_i'^*}$.

case 2 : $\mu_j^* = \mu_j, \mu_j' \neq \mu_j, \mu_j'^* \neq \mu_j$

In this case, the induction of $V_{\mu_j, \#}^+$ to $\mathcal{H}^\#$ yields $(V_{\mu_j, \#}^+)^{\mathcal{H}^\#} \cong V_{\mu_j, \#}^+ \oplus V_{\mu_j', \#}^+$ by Proposition 7.7 (1) and Theorem 2.7 (1). Moreover, the induction of $V_{\mu_j, \#}^+$ to $\mathcal{H}$ yields $(V_{\mu_j, \#}^+)^{\mathcal{H}} \cong V_{\mu_j}^+$ by Theorem 2.7 (2). Thus we obtain $(V_{\mu_j, \#}^+)^{\mathcal{H}} \cong V_{\mu_j}^+ \oplus V_{\mu_j'}^+$. Similarly, $(V_{\mu_j, \#}^-)^{\mathcal{H}} \cong V_{\mu_j}^- \oplus V_{\mu_j'}^-$ holds.

case 3 : $\nu_k^* \neq \nu_k, \nu_k' = \nu_k, \nu_k'^* \neq \nu_k$

Considering the induction of $V_{\nu_k, \#}^\dagger$ to $\mathcal{H}^\dagger$, we obtain $(V_{\nu_k, \#}^\dagger)^{\mathcal{H}} \cong V_{\nu_k} \oplus V_{\nu_k^*}$ in the same fashion as in the case 2.

case 4 : $\kappa_l^* \neq \kappa_l, \kappa_l' \neq \kappa_l, \kappa_l'^* = \kappa_l$

Considering the induction of $V_{\kappa_l, \#}^\dagger$ to $\mathcal{H}^\dagger$, we obtain $(V_{\kappa_l, \#}^\dagger)^{\mathcal{H}} \cong V_{\kappa_l} \oplus V_{\kappa_l'}$ in the same fashion as in the case 2.

case 5 : $\iota_m' = \iota_m^* = \iota_m'^* = \iota_m$

In this case, the induction of $V_{\iota_m, \#}^{++}$ to $\mathcal{H}^\#$ yields $(V_{\iota_m, \#}^{++})^{\mathcal{H}^\#} \cong V_{\iota_m, \#}^{++}$ by Corollary 7.8 and Theorem 2.7 (2). Moreover, the induction of $V_{\iota_m, \#}^{++}$ to $\mathcal{H}$ yields $(V_{\iota_m, \#}^{++})^{\mathcal{H}} \cong V_{\iota_m}^{++}$ by Theorem 2.7 (2). Thus we obtain $(V_{\iota_m, \#}^{++})^{\mathcal{H}} \cong V_{\iota_m}^{++}$. Similarly, we get $(V_{\iota_m, \#}^{+-})^{\mathcal{H}} \cong (V_{\iota_m, \#}^{-+})^{\mathcal{H}} \cong (V_{\iota_m, \#}^{--})^{\mathcal{H}} \cong V_{\iota_m}^{--}$.

Those induced representations are mutually non-isomorphic, hence we can conclude that the elements of $\text{Irr}(\mathcal{H}_{\bar{K}_1, B_n}(q)^\dagger)$ are mutually inequivalent.

Finally, we shall show the semisimplicity of $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger$. We define a map π_n to be

$$\pi_n : x \in \mathcal{H}^\dagger \longmapsto \bigoplus_{\pi \in \text{Irr}(\mathcal{H}^\dagger)} \pi(x)$$

Then, by theorems of Burnside and Frobenius-Schur, $\mathcal{H}^\dagger$ has a quotient $\pi_n(\mathcal{H}^\dagger)$ isomorphic to the semisimple algebra

$$\bigoplus_{\pi \in \text{Irr}(\mathcal{H}^\dagger)} \text{End}_{\bar{K}_1} \bar{K}_1^{\deg \pi} \cong \bigoplus_{\pi \in \text{Irr}(\mathcal{H}^\dagger)} \text{Mat}(\deg \pi, \bar{K}_1).$$

We claim that this semisimple algebra has dimension $2^{n-2}n!$. On the other hand, $\dim_{\bar{K}_1} \mathcal{H}^\dagger = 2^{n-2}n!$. Therefore we can conclude that $\text{Irr}(\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger)$ is a basic set of irreducible representations and that $\mathcal{H}_{\bar{K}_1, B_n}(1, q)^\dagger$ is semisimple. $\square$

References

- [1] S. Ariki and K. Koike, A Hecke algebra of $(\mathbb{Z}/r\mathbb{Z}) \wr \mathfrak{S}_n$ and construction of its irreducible representations, *Adv. Math.* **106** (1994), 216–243.
- [2] C. W. Curtis and I. Reiner, “Methods of Representation Theory,” Vol.1, John Wiley & Sons, 1981.
- [3] M. Geck and G. Pfeiffer, “Characters of finite Coxeter groups and Iwahori-Hecke algebras”, Oxford university press, 2000.
- [4] P. N. Hoefsmit, Representation of Hecke algebras of finite groups with BN-pairs of classical type, *Thesis* (1974), University of British Columbia.
- [5] H. Mitsuhashi, The q -analogue of the alternating group and its representations, *J. Algebra.* **240** (2001), 535–558.
- [6] H. Mitsuhashi, $\mathbb{Z}_2$ -graded Clifford system in the Hecke algebra of type B_n , *J. Algebra* **264**, (2003), 231–250.
- [7] C. Nastasescu and F. V. Oystaeyen, Methods of Graded Rings, Lecture Notes in Math., No. 1836, Springer-Verlag, Berlin, 2004.